

Nash equilibrium payoffs for stochastic differential games with reflection

Qian LIN *

Abstract In this paper, we investigate Nash equilibrium payoffs for nonzero-sum stochastic differential games with reflection. We obtain an existence theorem and a characterization theorem of Nash equilibrium payoffs for nonzero-sum stochastic differential games with nonlinear cost functionals defined by doubly controlled reflected backward stochastic differential equations.

Keywords Nash equilibrium payoff · Stochastic differential games · Backward stochastic differential equations · Dynamic programming principle

1 Introduction

In this paper we study Nash equilibrium payoffs for nonzero-sum stochastic differential games whose cost functionals are defined by doubly controlled reflected backward stochastic differential equations (RBSDEs, for short). Fleming and Souganidis [1] were the first in a rigorous way to study zero-sum stochastic differential games. Since the pioneering work of Fleming and Souganidis [1], stochastic differential games have been investigated by many authors. Recently, Buckdahn and Li [2] generalized the results of Fleming and Souganidis [1] by using a Girsanov transformation argument and a backward stochastic differential equation (BSDE, for short) approach. The reader interested in this topic can be referred to Buckdahn, Cardaliaguet and Quincampoix [3], Buckdahn and Li [2], Fleming and Souganidis [1] and the references therein.

El Karoui, Kapoudjian, Pardoux, Peng and Quenez [4] introduced RBSDEs. By virtue of RBSDEs they gave a probabilistic representation for the solution of an obstacle problem for a nonlinear parabolic partial differential equation. This kind of RBSDEs also has many applications in finance, stochastic differential games and stochastic optimal control problem. In [5], El Karoui, Pardoux and Quenez showed that the price of an American option corresponds to the solution of a RBSDE. Buckdahn and Li [13] considered zero-sum stochastic differential games with reflection. Wu and Yu [6] studied one kind of stochastic recursive optimal control problem with the obstacle constraint for the cost functional defined by a RBSDE.

Buckdahn, Cardaliaguet and Rainer [7] studied Nash equilibrium payoffs for stochastic differential games. Recently, Lin [8, 9] studied Nash equilibrium payoffs for stochastic differential games whose cost functionals are defined by doubly controlled BSDEs. Lin [8, 9] generalize the earlier result by Buckdahn, Cardaliaguet and Rainer [7]. In [8, 9], the admissible control processes can depend on events occurring before the beginning of the stochastic differential game, thus, the cost functionals are not necessarily deterministic. Moreover, the cost functionals are defined with the help of BSDEs, and thus they are nonlinear. The objective of this paper is to investigate Nash equilibrium payoffs for nonzero-sum stochastic differential games with reflection. However, different from the earlier results by Buckdahn, Cardaliaguet and Rainer [7] and Lin [8, 9], we shall study Nash equilibrium payoffs for stochastic differential games with general running cost functionals, which are defined with the help of RBSDEs. In comparison with Buckdahn and Li [13] we shall study nonzero-sum stochastic differential games of the type of "strat-

*Q. Lin

Laboratoire de Mathématiques, CNRS UMR 6205, Université de Bretagne Occidentale 6, avenue Victor Le Gorgeu, CS 93837, 29238 Brest cedex 3, France.
e-mail: Qian.Lin@univ-brest.fr

egy against strategy", while Buckdahn and Li [13] considered the games of the type "strategy against control". The related works see [10], [11] and [12].

The paper is organized as follows. In Section 2, we introduce some notations and present some preliminary results concerning reflected backward stochastic differential equations, which we will need in what follows. In Section 3, we introduce nonzero-sum stochastic differential games with reflection and obtain the associated dynamic programming principle. In Section 4 we give a probabilistic interpretation of systems of Isaacs equations with obstacle. In Section 5, we obtain the main results of this paper, i.e., an existence theorem and a characterization theorem of Nash equilibrium payoffs for nonzero-sum stochastic differential games with reflection.

2 Preliminaries

Let $B = \{B_t, 0 \leq t \leq T\}$ be a d -dimensional standard Brownian motion defined on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. The filtration $\mathbb{F} = \{\mathcal{F}_t, 0 \leq t \leq T\}$ is generated by B and augmented by all $\mathbb{P}$ -null sets, i.e.,

$$\mathcal{F}_t = \sigma\{B_r, 0 \leq r \leq t\} \vee \mathcal{N}_{\mathbb{P}},$$

where $\mathcal{N}_{\mathbb{P}}$ is the set of all $\mathbb{P}$ -null sets. Let us introduce some spaces:

- $L^2(\Omega, \mathcal{F}_T, \mathbb{P}; \mathbb{R}^n) = \left\{ \xi \mid \xi : \Omega \rightarrow \mathbb{R}^n \text{ is an } \mathcal{F}_T\text{-measurable random variable} \right.$
 $\left. \text{such that } \mathbb{E}[|\xi|^2] < +\infty \right\},$
- $S^2(0, T; \mathbb{R}) = \left\{ \varphi \mid \varphi : \Omega \times [0, T] \rightarrow \mathbb{R} \text{ is a predictable process such that} \right.$
 $\left. \mathbb{E}\left[\sup_{0 \leq t \leq T} |\varphi_t|^2\right] < +\infty \right\},$
- $\mathcal{H}^2(0, T; \mathbb{R}^d) = \left\{ \varphi \mid \varphi : \Omega \times [0, T] \rightarrow \mathbb{R}^d \text{ is a predictable process such that} \right.$
 $\left. \mathbb{E} \int_0^T |\varphi_t|^2 dt < +\infty \right\}.$

We consider the following one barrier reflected BSDE with data (f, ξ, S) :

$$\begin{cases} Y_t = \xi + \int_t^T f(s, Y_s, Z_s) ds + K_T - K_t - \int_t^T Z_s dW_s, \\ Y_t \geq S_t, \quad t \in [0, T], \\ K_0 = 0, \quad \int_0^T (Y_t - S_t) dK_t = 0, \end{cases} \quad (2.1)$$

where $\{K_t\}$ is an adapted, continuous and increasing process, $f : \Omega \times [0, T] \times \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{R}$ and we make the following assumptions:

(H2.1) $f(\cdot, 0, 0) \in \mathcal{H}^2(0, T; \mathbb{R})$,

(H2.2) There exists some constant $L > 0$ such that for all $y, y' \in \mathbb{R}$ and $z, z' \in \mathbb{R}^d$,

$$|f(t, y, z) - f(t, y', z')| \leq L(|y - y'| + |z - z'|),$$

(H2.3) $\{S_t\}_{t \in [0, T]}$ is a continuous process such that $\{S_t\}_{0 \leq t \leq T} \in S^2(0, T; \mathbb{R})$.

The following the existence and uniqueness theorem for solutions of equation (2.1) was established in [4].

Lemma 2.1. *Under the assumptions (H2.1)-(H2.3), if $\xi \in L^2(\Omega, \mathcal{F}_T, \mathbb{P}; \mathbb{R})$ and $S_T \leq \xi$, then equation (2.1) has a unique solution (Y, Z, K) .*

We refer to [4] and [6] for the following two estimates.

Lemma 2.2. *Let assumptions (H2.1)-(H2.3) hold and let (Y, Z, K) be the solution of the reflected BSDE (2.1) with data (ξ, g, S) . Then there exists a positive constant C such that*

$$\mathbb{E}\left[\sup_{t \leq s \leq T} Y_s^2 + \int_t^T |Z_s|^2 + |K_T - K_t|^2 \middle| \mathcal{F}_t\right] \leq C\mathbb{E}\left[\xi^2 + \left(\int_t^T g(s, 0, 0)ds\right)^2 + \sup_{t \leq s \leq T} S_s^2 \middle| \mathcal{F}_t\right].$$

Lemma 2.3. *We suppose that (ξ, g, S) and (ξ', g', S') satisfy assumptions (H2.1)-(H2.3). Let (Y, Z, K) and (Y', Z', K') be the solutions of the reflected BSDEs (2.1) with data (ξ, g, S) and (ξ', g', S') , respectively. We let*

$$\begin{aligned} \Delta\xi &= \xi - \xi', & \Delta g &= g - g', & \Delta S &= S - S', \\ \Delta Y &= Y - Y', & \Delta Z &= Z - Z', & \Delta K &= K - K'. \end{aligned}$$

Then there exists a constant C such that

$$\begin{aligned} & \mathbb{E}\left[\sup_{t \leq s \leq T} |\Delta Y_s|^2 + \int_t^T |\Delta Z_s|^2 ds + |\Delta K_T - \Delta K_t|^2 \middle| \mathcal{F}_t\right] \\ & \leq C\mathbb{E}\left[|\Delta\xi|^2 + \left(\int_t^T |\Delta g(s, Y_s, Z_s)|ds\right)^2 \middle| \mathcal{F}_t\right] + C\left(\mathbb{E}\left[\sup_{t \leq s \leq T} |\Delta S_s|^2 \middle| \mathcal{F}_t\right]\right)^{1/2} \Psi_{t,T}^{1/2}, \end{aligned}$$

where

$$\begin{aligned} \Psi_{t,T} &= \mathbb{E}\left[|\xi|^2 + \left(\int_t^T |g(s, 0, 0)|ds\right)^2 + \sup_{t \leq s \leq T} |S_s|^2 \right. \\ & \quad \left. + |\xi'|^2 + \left(\int_t^T |g'(s, 0, 0)|ds\right)^2 + \sup_{t \leq s \leq T} |S'_s|^2 \middle| \mathcal{F}_t\right]. \end{aligned}$$

We also need the following lemma. For its proof, the interested reader can refer to [4] and [14] for more details.

Lemma 2.4. *Let us denote by (Y^1, Z^1, K^1) and (Y^2, Z^2, K^2) the solutions of BSDEs with data (f^1, ξ^1, S^1) and (f^2, ξ^2, S^2) , respectively. If $\xi^1, \xi^2 \in L^2(\Omega, \mathcal{F}_T, \mathbb{P}; \mathbb{R})$, S^1 and S^2 satisfy (H2.3), and f^1 and f^2 satisfy the assumptions (H2.1) and (H2.2), and the following holds*

- (i) $\xi^1 \leq \xi^2$, $\mathbb{P}$ -a.s.,
- (ii) $f^1(t, y_t^2, z_t^2) \leq f^2(t, y_t^2, z_t^2)$, $dt d\mathbb{P}$ -a.e
- (iii) $S^1 \leq S^2$, $\mathbb{P}$ -a.s.

Then, we have $Y_t^1 \leq Y_t^2$, a.s., for all $t \in [0, T]$. Moreover, if

- (iv) $f^1(t, y, z) \leq f^2(t, y, z)$, $(t, y, z) \in [0, T] \times \mathbb{R} \times \mathbb{R}^d$, $dt d\mathbb{P}$ -a.e
- (v) $S^1 = S^2$, $\mathbb{P}$ -a.s.

Then, $K_t^1 \leq K_t^2$, $\mathbb{P}$ -a.s., for all $t \in [0, T]$, and $\{K^1 - K^2\}_{t \in [0, T]}$ is a decreasing process.

3 Nonzero-sum stochastic differential games with reflection and associated dynamic programming principle

In what follows, we assume that U and V are two compact metric spaces. The space U is considered as the control state space of the first player, and V as that of the second one. We denote the associated sets of admissible controls by $\mathcal{U}$ and $\mathcal{V}$, respectively. The set $\mathcal{U}$ is formed by all U -valued $\mathbb{F}$ -progressively measurable processes, and $\mathcal{V}$ is the set of all V -valued $\mathbb{F}$ -progressively measurable processes.

For given admissible controls $u(\cdot) \in \mathcal{U}$ and $v(\cdot) \in \mathcal{V}$, let us consider the following control system: for $t \in [0, T]$,

$$\begin{cases} dX_s^{t,x;u,v} &= b(s, X_s^{t,x;u,v}, u_s, v_s)ds + \sigma(s, X_s^{t,x;u,v}, u_s, v_s)dB_s, & s \in [t, T], \\ X_t^{t,x;u,v} &= x \in \mathbb{R}^n, \end{cases} \quad (3.1)$$

where

$$b : [0, T] \times \mathbb{R}^n \times U \times V \rightarrow \mathbb{R}^n, \quad \sigma : [0, T] \times \mathbb{R}^n \times U \times V \rightarrow \mathbb{R}^{n \times d}.$$

We make the following assumptions:

(H3.1) For all $x \in \mathbb{R}^n$, $b(\cdot, x, \cdot, \cdot)$ and $\sigma(\cdot, x, \cdot, \cdot)$ are continuous in (t, u, v) .

(H3.2) There exists a positive constant L such that, for all $t \in [0, T]$, $x, x' \in \mathbb{R}^n$, $u \in U, v \in V$,

$$|b(t, x, u, v) - b(t, x', u, v)| + |\sigma(t, x, u, v) - \sigma(t, x', u, v)| \leq L|x - x'|.$$

Under the above assumptions, for any $u(\cdot) \in \mathcal{U}$ and $v(\cdot) \in \mathcal{V}$, the control system (3.1) has a unique strong solution $\{X_s^{t,x;u,v}, 0 \leq t \leq s \leq T\}$, and we also have the following standard estimates for solutions.

Lemma 3.1. *For all $p \geq 2$, there exists a positive constant C_p such that, for all $t \in [0, T]$, $x, x' \in \mathbb{R}^n$, $u(\cdot) \in \mathcal{U}$ and $v(\cdot) \in \mathcal{V}$,*

$$\begin{aligned} \mathbb{E} \left[\sup_{t \leq s \leq T} |X_s^{t,x;u,v}|^p \middle| \mathcal{F}_t \right] &\leq C_p(1 + |x|^p), \quad \mathbb{P} - a.s., \\ \mathbb{E} \left[\sup_{t \leq s \leq T} |X_s^{t,x;u,v} - X_s^{t,x';u,v}|^p \middle| \mathcal{F}_t \right] &\leq C_p|x - x'|^p, \quad \mathbb{P} - a.s., \end{aligned}$$

where the constant C_p only depends on p , the Lipschitz constant and the linear growth of b and σ .

For given admissible controls $u(\cdot) \in \mathcal{U}$ and $v(\cdot) \in \mathcal{V}$, let us consider the following doubly controlled reflected BSDE:

$$\begin{cases} {}^jY_s^{t,x;u,v} = \Phi_j(X_T^{t,x;u,v}) + \int_s^T f_j(r, X_r^{t,x;u,v}, {}^jY_r^{t,x;u,v}, {}^jZ_r^{t,x;u,v}, u_r, v_r)dr \\ \quad + {}^jK_T^{t,x;u,v} - {}^jK_s^{t,x;u,v} - \int_s^T {}^jZ_r^{t,x;u,v}dB_r, & s \in [t, T], \\ {}^jY_s^{t,x;u,v} \geq h_j(s, X_s^{t,x;u,v}), & s \in [t, T], \\ {}^jK_t^{t,x;u,v} = 0, \quad \int_t^T ({}^jY_r^{t,x;u,v} - h_j(r, X_r^{t,x;u,v}))d {}^jK_r^{t,x;u,v} = 0. \end{cases} \quad (3.2)$$

where $X^{t,x;u,v}$ is introduced in equation (3.1) and

$$\begin{aligned} \Phi_j &= \Phi_j(x) : \mathbb{R}^n \rightarrow \mathbb{R}, \quad h_j = h_j(t, x) : [0, T] \times \mathbb{R}^n \rightarrow \mathbb{R}, \\ f_j &= f_j(t, x, y, z, u, v) : [0, T] \times \mathbb{R}^n \times \mathbb{R} \times \mathbb{R}^d \times U \times V \rightarrow \mathbb{R}. \end{aligned}$$

We make the following assumptions:

(H3.3) There exists a positive constant L such that, for all $t \in [0, T]$, $x, x' \in \mathbb{R}^n$, $y, y' \in \mathbb{R}$, $z, z' \in \mathbb{R}^d$, $u \in U$ and $v \in V$,

$$\begin{aligned} &|f_j(t, x, y, z, u, v) - f_j(t, x', y', z', u, v)| + |\Phi_j(x) - \Phi_j(x')| + |h_j(t, x) - h_j(t, x')| \\ &\leq L(|x - x'| + |y - y'| + |z - z'|). \end{aligned}$$

(H3.4) For all $(x, y, z) \in \mathbb{R}^n \times \mathbb{R} \times \mathbb{R}^d$, $f_j(\cdot, x, y, z, \cdot, \cdot)$ is continuous in (t, u, v) , and there exists a positive constant L such that, for all $t, s \in [0, T]$, $x \in \mathbb{R}^n$,

$$|h_j(t, x) - h_j(s, x)| \leq L|t - s|.$$

Under the assumption (H3.3), from [4] we know that equation (3.2) admits a unique solution.

For given control processes $u(\cdot) \in U$ and $v(\cdot) \in V$, let us introduce now the associated cost functional for player j , $j = 1, 2$,

$$J_j(t, x; u, v) := {}^j Y_s^{t, x; u, v} \Big|_{s=t}, \quad (t, x) \in [0, T] \times \mathbb{R}^n.$$

From Buckdahn and Li [13] we have the following estimates for solutions.

Proposition 3.2. *Under the assumption (H3.1)-(H3.3), there exists a positive constant C such that, for all $t \in [0, T]$, $u(\cdot) \in \mathcal{U}$ and $v(\cdot) \in \mathcal{V}$, $x, x' \in \mathbb{R}^n$,*

$$\begin{aligned} |{}^j Y_t^{t, x; u, v}| &\leq C(1 + |x|), \quad \mathbb{P} - a.s., \\ |{}^j Y_t^{t, x; u, v} - {}^j Y_t^{t, x'; u, v}| &\leq C|x - x'|, \quad \mathbb{P} - a.s. \end{aligned}$$

We now recall the definition of admissible controls and NAD strategies, which were introduced in [8].

Definition 3.3. *The space $\mathcal{U}_{t, T}$ (resp., $\mathcal{V}_{t, T}$) of admissible controls for 1^{th} player (resp., 2^{th}) on the interval $[t, T]$ is defined as the space of all processes $\{u_r\}_{r \in [t, T]}$ (resp., $\{v_r\}_{r \in [t, T]}$), which are $\mathbb{F}$ -progressively measurable and take values in U (resp., V).*

Definition 3.4. *A nonanticipating strategy with delay (NAD strategy) for 1^{th} player is a measurable mapping $\alpha : \mathcal{V}_{t, T} \rightarrow \mathcal{U}_{t, T}$, which satisfies the following properties:*

1) α is a nonanticipative strategy, i.e., for every $\mathbb{F}$ -stopping time $\tau : \Omega \rightarrow [t, T]$, and for $v_1, v_2 \in \mathcal{V}_{t, T}$ with $v_1 = v_2$ on $[[t, \tau]]$, it holds $\alpha(v_1) = \alpha(v_2)$ on $[[t, \tau]]$. (Recall that $[[t, \tau]] = \{(s, \omega) \in [t, T] \times \Omega, t \leq s \leq \tau(\omega)\}$).

2) α is a strategy with delay, i.e., for all $v \in \mathcal{V}_{t, T}$, there exists an increasing sequence of stopping times $\{S_n(v)\}_{n \geq 1}$ with

i) $t = S_0(v) \leq S_1(v) \leq \dots \leq S_n(v) \leq \dots \leq T$,

ii) $\bigcup_{n \geq 1} \{S_n(v) = T\} = \Omega$, $\mathbb{P}$ -a.s.,

such that, for all $n \geq 1$ and $v, v' \in \mathcal{V}_{t, T}$, $\Gamma \in \mathcal{F}_t$, it holds: if $v = v'$ on $[[t, S_{n-1}(v)]] \cap (\Gamma \times [t, T])$, then

iii) $S_l(v) = S_l(v')$, on Γ , $1 \leq l \leq n$,

iv) $\alpha(v) = \alpha(v')$, on $[[t, S_n(v)]] \cap (\Gamma \times [t, T])$.

We denote the set of all NAD strategies for 1^{th} player for games over the time interval $[t, T]$ by $\mathcal{A}_{t, T}$. The set of all NAD strategies $\beta : \mathcal{U}_{t, T} \rightarrow \mathcal{V}_{t, T}$ for 2^{th} player for games over the time interval $[t, T]$ is defined in a symmetrical way and we denote it by $\mathcal{B}_{t, T}$.

NAD strategy allows us to put stochastic differential games under normal form. The following lemma was established in [8].

Lemma 3.5. *If $(\alpha, \beta) \in \mathcal{A}_{t, T} \times \mathcal{B}_{t, T}$, then there exists a unique couple of admissible control $(u, v) \in \mathcal{U}_{t, T} \times \mathcal{V}_{t, T}$ such that*

$$\alpha(v) = u, \quad \beta(u) = v.$$

If $(\alpha, \beta) \in \mathcal{A}_{t, T} \times \mathcal{B}_{t, T}$, then from Lemma 3.5 we have a unique couple $(u, v) \in \mathcal{U}_{t, T} \times \mathcal{V}_{t, T}$ such that $(\alpha(v), \beta(u)) = (u, v)$. Then let us put $J_j(t, x; \alpha, \beta) = J_j(t, x; u, v)$. Therefore, let us define the lower and the upper value functions W_j and U_j , resp., of the game: for all $(t, x) \in [0, T] \times \mathbb{R}^n$, we put

$$W_j(t, x) := \operatorname{esssup}_{\alpha \in \mathcal{A}_{t, T}} \operatorname{essinf}_{\beta \in \mathcal{B}_{t, T}} J_j(t, x; \alpha, \beta),$$

and

$$U_j(t, x) := \operatorname{essinf}_{\beta \in \mathcal{B}_{t, T}} \operatorname{esssup}_{\alpha \in \mathcal{A}_{t, T}} J_j(t, x; \alpha, \beta).$$

Under assumptions (H3.1)–(H3.3) we see that $W_j(t, x)$ and $U_j(t, x)$ are random variables. But using the arguments in [3] and [9], we have the following proposition.

Proposition 3.6. *Under assumptions (H3.1)–(H3.3), for all $(t, x) \in [0, T] \times \mathbb{R}^n$, the value functions $W_j(t, x)$ and $U_j(t, x)$ are deterministic functions.*

Let us now recall the definition of stochastic backward semigroups, which was first introduced by Peng [15] to study stochastic optimal control problem. For a given initial condition (t, x) , a positive number $\delta \leq T - t$, for admissible control processes $u(\cdot) \in \mathcal{U}_{t, t+\delta}$ and $v(\cdot) \in \mathcal{V}_{t, t+\delta}$, and a real-valued random variable $\eta \in L^2(\Omega, \mathcal{F}_{t+\delta}, \mathbb{P}; \mathbb{R})$ such that $\eta \geq h_j(t + \delta, X_{t+\delta}^{t, x; u, v})$, we define

$${}^j G_{t, t+\delta}^{t, x; u, v}[\eta] := {}^j \bar{Y}_t^{t, x; u, v},$$

where $({}^j \bar{Y}_s^{t, x; u, v}, {}^j \bar{Z}_s^{t, x; u, v}, {}^j \bar{K}_s^{t, x; u, v})$ is the unique solution of the following reflected BSDE over the time interval $[t, t + \delta]$:

$$\begin{cases} {}^j \bar{Y}_s^{t, x; u, v} = \eta + \int_s^{t+\delta} f_j(r, X_r^{t, x; u, v}, {}^j \bar{Y}_r^{t, x; u, v}, {}^j \bar{Z}_r^{t, x; u, v}, u_r, v_r) dr \\ \quad + {}^j \bar{K}_{t+\delta}^{t, x; u, v} - {}^j \bar{K}_s^{t, x; u, v} - \int_s^{t+\delta} {}^j \bar{Z}_r^{t, x; u, v} dB_r, \quad s \in [t, t + \delta], \\ {}^j \bar{Y}_s^{t, x; u, v} \geq h_j(s, X_s^{t, x; u, v}), \quad s \in [t, t + \delta], \\ {}^j \bar{K}_t^{t, x; u, v} = 0, \quad \int_t^{t+\delta} ({}^j \bar{Y}_r^{t, x; u, v} - h_j(r, X_r^{t, x; u, v})) d{}^j \bar{K}_r^{t, x; u, v} = 0. \end{cases}$$

and $X^{t, x; u, v}$ is the unique solution of equation (3.1).

For $(t, x) \in [0, T] \times \mathbb{R}^n$, $(u, v) \in \mathcal{U}_{t, T} \times \mathcal{V}_{t, T}$, $0 \leq \delta \leq T - t$, $j = 1, 2$, we have

$$\begin{aligned} J_j(t, x; u, v) &= {}^j G_{t, T}^{t, x; u, v}[\Phi_j(X_T^{t, x; u, v})] = {}^j G_{t, t+\delta}^{t, x; u, v}[{}^j Y_{t+\delta}^{t, x; u, v}] \\ &= {}^j G_{t, t+\delta}^{t, x; u, v}[J_j(t + \delta, X_{t+\delta}^{t, x; u, v}, u, v)], \end{aligned}$$

Remark 3.7. *We consider a special case of f_j . If f_j is independent of (y, z) , then we have*

$$J_j(t, x; u, v) = {}^j G_{t, t+\delta}^{t, x; u, v}[\eta] = \mathbb{E}\left[\eta + \int_t^{t+\delta} f_j(r, X_r^{t, x; u, v}, u_r, v_r) dr + {}^j \bar{K}_{t+\delta}^{t, x; u, v} \mid \mathcal{F}_t\right].$$

Proposition 3.8. *Under the assumptions (H3.1)–(H3.3) we have the following dynamic programming principle: for all $0 < \delta \leq T - t$, $x \in \mathbb{R}^n$,*

$$W_j(t, x) = \operatorname{esssup}_{\alpha \in \mathcal{A}_{t, t+\delta}} \operatorname{essinf}_{\beta \in \mathcal{B}_{t, t+\delta}} {}^j G_{t, t+\delta}^{t, x; \alpha, \beta}[W_j(t + \delta, X_{t+\delta}^{t, x; \alpha, \beta})],$$

and

$$U_j(t, x) = \operatorname{essinf}_{\beta \in \mathcal{B}_{t, t+\delta}} \operatorname{esssup}_{\alpha \in \mathcal{A}_{t, t+\delta}} {}^j G_{t, t+\delta}^{t, x; \alpha, \beta}[U_j(t + \delta, X_{t+\delta}^{t, x; \alpha, \beta})].$$

The proof of the above proposition is similar to [3] and [9], we omit the proof here.

Proposition 3.9. *Under the assumptions (H3.1)–(H3.4), there exists a positive constant C such that, for all $t, t' \in [0, T]$ and $x, x' \in \mathbb{R}^n$, we have*

(i) $W_j(t, x)$ is $\frac{1}{2}$ -Hölder continuous in t :

$$|W_j(t, x) - W_j(t', x)| \leq C(1 + |x|)|t - t'|^{\frac{1}{2}};$$

(ii) $|W_j(t, x) - W_j(t, x')| \leq C|x - x'|$.

The same properties hold true for the function U_j .

By means of the standard arguments and Proposition 3.8 we can easily get the above proposition. The proof of the above proposition is omitted here.

4 Probabilistic interpretation of systems of Isaacs equations with obstacle

The objective of this section is to give a probabilistic interpretation of systems of Isaacs equations with obstacle, and show that W_j and U_j introduced in Section 3, are the viscosity solutions of the following Isaacs equations with obstacle, for $(t, x) \in [0, T] \times \mathbb{R}^n$,

$$\begin{cases} \min \left\{ W_j(t, x) - h_j(t, x), -\frac{\partial}{\partial t} W_j(t, x) - H_j^-(t, x, W_j(t, x), DW_j(t, x), D^2 W_j(t, x)) \right\} = 0, \\ W_j(T, x) = \Phi_j(x), \end{cases} \quad (4.1)$$

and

$$\begin{cases} \min \left\{ U_j(t, x) - h_j(t, x), -\frac{\partial}{\partial t} U_j(t, x) - H_j^+(t, x, U_j(t, x), DU_j(t, x), D^2 U_j(t, x)) \right\} = 0, \\ U_j(T, x) = \Phi_j(x), \end{cases} \quad (4.2)$$

respectively, where

$$\begin{aligned} H_j(t, x, y, p, A, u, v) &= \frac{1}{2} \text{tr}(\sigma \sigma^T(t, x, u, v) A_j) + p b(t, x, u, v) \\ &\quad + f_j(t, x, y, p, \sigma(t, x, u, v), u, v), \end{aligned}$$

$$(t, x, y, p, A, u, v) \in [0, T] \times \mathbb{R}^n \times \mathbb{R} \times \mathbb{R}^d \times \mathbb{S}^d \times U \times V,$$

$$H_j^-(t, x, y, p, A) = \sup_{u \in U} \inf_{v \in V} H_j(t, x, y, p, A, u, v)$$

and

$$H_j^+(t, x, y, p, A) = \inf_{v \in V} \sup_{u \in U} H_j(t, x, y, p, A, u, v)$$

We denote by $C_{l,b}^3([0, T] \times \mathbb{R}^n)$ the set of real-valued functions which are continuously differentiable up to the third order and whose derivatives of order from 1 to 3 are bounded. Let us recall the definition of a viscosity solution of (4.1). The definition of a viscosity solution of (4.2) can be defined in a similar way. The interested reader is referred to Crandall, Ishii and Lions [16] for more details.

Definition 4.1. Let $w \in C([0, T] \times \mathbb{R}^n; \mathbb{R})$ be a function. It is called

(i) a viscosity subsolution of (4.1) if

$$w_j(T, x) \leq \Phi_j(x), \text{ for all } x \in \mathbb{R}^n,$$

and if for all functions $\varphi \in C_{l,b}^3([0, T] \times \mathbb{R}^n)$, $j = 1, 2$, and $(t, x) \in [0, T] \times \mathbb{R}^n$ such that $w_j - \varphi$ attains a local maximum at (t, x) ,

$$\min \left\{ w_j(t, x) - h_j(t, x), -\frac{\partial}{\partial t} \varphi_j(t, x) - H_j^-(t, x, w_j(t, x), D\varphi_j(t, x), D^2 \varphi_j(t, x)) \right\} \leq 0,$$

(ii) a viscosity supsolution of (4.1) if

$$w_i(T, x) \geq \Phi_i(x), \text{ for all } j = 1, 2, x \in \mathbb{R}^n,$$

and if for all functions $\varphi \in C_{l,b}^3([0, T] \times \mathbb{R}^n)$, $j = 1, 2$, and $(t, x) \in [0, T] \times \mathbb{R}^n$ such that $w_j - \varphi$ attains a local minimum at (t, x) ,

$$\min \left\{ w_j(t, x) - h_j(t, x), -\frac{\partial}{\partial t} w_j(t, x) - H_j^-(t, x, w_j(t, x), Dw_j(t, x), D^2 w_j(t, x)) \right\} \geq 0,$$

(iii) a viscosity solution of (4.1) if it is both a viscosity subsolution and a supersolution of (4.1).

Theorem 4.2. *Under the assumptions (H3.1)–(H3.3), the function W_j (resp., U_j) is a viscosity solution of the system (4.1) (resp., (4.2)).*

Proof. For arbitrarily fixed $j = 1, 2$, we only prove that the function W_j is a viscosity solution of the system (4.1). Let us consider the following BSDE, for $(t, x) \in [0, T] \times \mathbb{R}^n$, $m \in \mathbb{N}$,

$$\begin{aligned} {}^{j,m}Y_s^{t,x;u,v} &= \Phi_j(X_T^{t,x;u,v}) + \int_s^T f_j(r, X_r^{t,x;u,v}, {}^{j,m}Y_r^{t,x;u,v}, {}^{j,m}Z_r^{t,x;u,v}, u_r, v_r) dr \\ &\quad + \int_s^T m({}^{j,m}Y_r^{t,x;u,v} - h_j(r, X_r^{t,x;u,v}))^- dr - \int_s^T {}^{j,m}Z_r^{t,x;u,v} dB_r, \quad s \in [t, T], \end{aligned}$$

and define

$$J_{j,m}(t, x; u, v) := {}^{j,m}Y_s^{t,x;u,v} \Big|_{s=t}.$$

From Lemma 3.5 it follows that there exists a unique couple $(u, v) \in \mathcal{U}_{t,T} \times \mathcal{V}_{t,T}$ such that $(\alpha(v), \beta(u)) = (u, v)$ thanks to $(\alpha, \beta) \in \mathcal{A}_{t,T} \times \mathcal{B}_{t,T}$. Hence, we can define $J_{j,m}(t, x; \alpha, \beta) = J_{j,m}(t, x; u, v)$, and we put

$$W_{j,m}(t, x) := \operatorname{esssup}_{\alpha \in \mathcal{A}_{t,T}} \operatorname{essinf}_{\beta \in \mathcal{B}_{t,T}} J_{j,m}(t, x; \alpha, \beta).$$

Using a similar argument to that in [3] we see that $W_{j,m}$ is of linear growth and is the unique viscosity solution of the following Isaacs equation:

$$\begin{cases} -\frac{\partial}{\partial t} W_{j,m}(t, x) - H_{j,m}^-(t, x, W_{j,m}(t, x), DW_{j,m}(t, x), D^2 W_{j,m}(t, x)) &= 0, \\ W_{j,m}(T, x) &= \Phi_j(x), \end{cases} \quad (4.3)$$

where for $(t, x, y, p, A, u, v) \in [0, T] \times \mathbb{R}^n \times \mathbb{R} \times \mathbb{R}^d \times \mathbb{S}^d \times U \times V$,

$$\begin{aligned} H_{j,m}^-(t, x, y, p, A) &= \sup_{u \in U} \inf_{v \in V} \left\{ \frac{1}{2} \operatorname{tr}(\sigma \sigma^T(t, x, u, v) A) + pb(t, x, u, v) \right. \\ &\quad \left. + f_j(t, x, y, p\sigma(t, x, u, v), u, v) + m(y - h(t, x))^- \right\}. \end{aligned}$$

Using an argument similar to that in Buckdahn and Li [13] we have the following lemma.

Lemma 4.3. *For every $(t, x) \in [0, T] \times \mathbb{R}^n$ and $m \geq 1$, the following holds:*

$$W_{j,m}(t, x) \leq W_{j,m+1}(t, x) \leq W_j(t, x),$$

and

$$\widetilde{W}_j(t, x) = \lim_{m \uparrow \infty} W_{j,m+1}(t, x),$$

is a viscosity supersolution of (4.1).

Let us admit for a moment the following lemma, and its proof will be given later.

Lemma 4.4. *For every $(t, x) \in [0, T] \times \mathbb{R}^n$, $W_j(t, x)$ is a viscosity subsolution of (4.1).*

From the above both lemmas and Theorem 4.7 it follows that $\widetilde{W}_j(t, x) = W_j(t, x)$. The desired result then follows. $\square$

Let us come to the **Proof of Lemma 4.4.**

Proof. Let $j = 1, 2$ be arbitrarily fixed, $(t, x) \in [0, T] \times \mathbb{R}^n$, and $\delta \in [0, T - t]$. For $\varphi \in C_{l,b}^3([0, T] \times \mathbb{R}^n)$, we define

$$\begin{aligned} F_j(s, x, y, z, u, v) &= \frac{\partial}{\partial s} \varphi(s, x) + \frac{1}{2} \text{tr}(\sigma \sigma^T(s, x, u, v) D^2 \varphi(s, x)) + D\varphi(s, x) b(s, x, u, v) \\ &\quad + f_j(s, x, y + \varphi(s, x), z + D\varphi(s, x) \cdot \sigma(s, x, u, v), u, v), \end{aligned}$$

where $(s, x, y, z, u, v) \in [0, T] \times \mathbb{R}^n \times \mathbb{R} \times \mathbb{R}^d \times U \times V$.

We consider the following BSDE:

$$\left\{ \begin{array}{l} Y_s^{1,u,v} = \int_s^{t+\delta} F_j(r, X_r^{t,x;u,v}, Y_r^{1,u,v}, Z_r^{1,u,v}, u_r, v_r) dr + K_{t+\delta}^{1,u,v} - K_s^{1,u,v} - \int_s^{t+\delta} Z_r^{1,u,v} dB_r, \\ Y_s^{1,u,v} \geq h_j(s, X_s^{t,x;u,v}) - \varphi(s, X_s^{t,x;u,v}), \quad s \in [t, t+\delta], \\ \int_t^{t+\delta} (Y_r^{1,u,v} - h_j(r, X_r^{t,x;u,v}) - \varphi(r, X_r^{t,x;u,v})) dK_r^{1,u,v} = 0, \\ K_t^{1,u,v} = 0, \end{array} \right. \quad (4.4)$$

where $X^{t,x;u,v}$ is the solution of (3.1).

Note that $F(s, x, y, h, z, u, v)$ is Lipschitz in (y, h, z) , uniformly in (s, x, u, v) , and there exists a positive constant C such that

$$F_j(s, x, 0, 0, 0, u, v) \leq C(1 + |x|^2), \quad (s, x, u, v) \in [0, T] \times \mathbb{R}^n \times U \times V.$$

Therefore, equation (4.4) has a unique solution $(Y^{1,u,v}, Z^{1,u,v}, K^{1,u,v})$. Moreover, we have the following lemma.

Lemma 4.5. *For every $s \in [t, t + \delta]$, we have*

$$Y_s^{1,u,v} = {}^j G_{s,t+\delta}^{t,x;u,v} [\varphi(t + \delta, X_{t+\delta}^{t,x;u,v})] - \varphi(s, X_s^{t,x;u,v}).$$

Let us also consider the following reflected BSDE:

$$\left\{ \begin{array}{l} Y_s^{2,u,v} = \int_s^{t+\delta} F_j(r, x, Y_r^{2,u,v}, Z_r^{2,u,v}, u_s, v_r) dr + K_{t+\delta}^{2,u,v} - K_s^{2,u,v} - \int_s^{t+\delta} Z_r^{2,u,v} dB_r, \\ Y_s^{2,u,v} \geq h_j(s, X_s^{t,x;u,v}) - \varphi(s, X_s^{t,x;u,v}), \quad s \in [t, t+\delta], \\ \int_t^{t+\delta} (Y_r^{2,u,v} - h_j(r, X_r^{t,x;u,v}) - \varphi(r, X_r^{t,x;u,v})) dK_r^{2,u,v} = 0, \\ K_t^{2,u,v} = 0. \end{array} \right. \quad (4.5)$$

Then we have the following lemma which estimates the difference of $Y_t^{1,u,v}$ and $Y_t^{2,u,v}$.

Lemma 4.6. *For $u \in \mathcal{U}_{t,t+\delta}$, $v \in \mathcal{V}_{t,t+\delta}$, there exists a positive constant C such that*

$$|Y_t^{1,u,v} - Y_t^{2,u,v}| \leq C\delta^{\frac{3}{2}}, \quad \mathbb{P} - a.s.,$$

where the constant C does not depend on δ , neither on the control processes u and v .

Using the standard arguments for BSDEs and Lemma 2.3 we can obtain the above two lemmas. We omit the proof here.

Let us prove that W_j is a viscosity subsolution of (4.1). For arbitrarily fixed $j = 1, 2$, we suppose that $W_j \leq \varphi$, $W_j(t, x) = \varphi(t, x)$ and $W_j(t, x) > h(t, x)$. Since W_j and h are continuous there exist $\delta_0 > 0$ and $\tau > 0$ such that

$$W_j(s, x') - h_j(s, x') \geq \tau, \quad \text{for } (s, x') \in [0, T] \times \mathbb{R}^n \text{ such that } |(t, x) - (s, x')| \leq \delta_0.$$

and we can choose φ such that

$$W_j(s, x') - \varphi(s, x') \leq -\tau, \quad \text{for } (s, x') \in [0, T] \times \mathbb{R}^n \text{ such that } |(t, x) - (s, x')| \geq \delta_0.$$

Consequently,

$$h_j(t, x) - \varphi(t, x) \leq -\tau, \quad \text{for } (t, x) \in [0, T] \times \mathbb{R}^n.$$

In what follows we shall prove that:

$$\inf_{v \in V} \sup_{u \in U} F_j(t, x, 0, 0, u, v) \geq 0.$$

We suppose that this is not true. Then, due to the continuity of F , there exist some $\theta > 0, v^* \in V$ and $0 < \delta' \leq T - t$ such that

$$\sup_{u \in U} F_j(s, x, 0, 0, u, v^*) \leq -\theta < 0, \quad (4.6)$$

for all $s \in [t, t + \delta']$.

By virtue of the Lemmas 2.4 and 3.5 we see that

$$\begin{aligned} Y_t^{1, \alpha, \beta} &= {}^j G_{t, t+\delta}^{t, x; \alpha, \beta} [\varphi(t + \delta, X_{t+\delta}^{t, x; \alpha, \beta})] - \varphi(s, X_s^{t, x; \alpha, \beta}) \\ &\geq {}^j G_{t, t+\delta}^{t, x; \alpha, \beta} [W_j(t + \delta, X_{t+\delta}^{t, x; \alpha, \beta})] - \varphi(t, x). \end{aligned}$$

On the other hand, from Proposition 3.8 it follows that

$$\text{esssup}_{\alpha \in \mathcal{A}_{t, t+\delta}} \text{essinf}_{\beta \in \mathcal{B}_{t, t+\delta}} {}^j G_{t, t+\delta}^{t, x; \alpha, \beta} [W_j(t + \delta, X_{t+\delta}^{t, x; \alpha, \beta})] - \varphi(t, x) = W_j(t, x) - \varphi(t, x) = 0.$$

Consequently,

$$\text{esssup}_{\alpha \in \mathcal{A}_{t, t+\delta}} \text{essinf}_{\beta \in \mathcal{B}_{t, t+\delta}} Y_t^{1, \alpha, \beta} \geq 0.$$

Let us define $\beta^*(u) = v^*$ on $[t, \tau^\delta] \times \mathcal{U}_{t, \tau^\delta}$. Then $\beta^* \in \mathcal{B}_{t, \tau^\delta}$. By Lemma 4.6 we have

$$\begin{aligned} 0 &\leq \text{esssup}_{\alpha \in \mathcal{A}_{t, t+\delta}} \text{essinf}_{\beta \in \mathcal{B}_{t, t+\delta}} Y_t^{1, \alpha, \beta} \\ &\leq \text{esssup}_{\alpha \in \mathcal{A}_{t, \tau^\delta}} Y_t^{1, \alpha, \beta^*} \\ &\leq \text{esssup}_{\alpha \in \mathcal{A}_{t, \tau^\delta}} Y_t^{2, \alpha, \beta^*} + C\delta^{\frac{3}{2}} \\ &= \text{esssup}_{\alpha \in \mathcal{A}_{t, \tau^\delta}} Y_t^{2, \alpha(v^*), v^*} + C\delta^{\frac{3}{2}} \\ &\leq \text{esssup}_{u \in \mathcal{U}_{t, \tau^\delta}} Y_t^{2, u, v^*} + C\delta^{\frac{3}{2}}, \end{aligned} \quad (4.7)$$

where we have used that $Y_t^{2, \alpha, \beta^*} = Y_t^{2, \alpha(v^*), v^*}$, since $\beta^*(\alpha(v^*)) = v^*$. Moreover, for all $\delta > 0, \varepsilon > 0$, we have the existence $u^\varepsilon \in \mathcal{U}_{t, \tau^\delta}$ such that

$$Y_t^{2, u^\varepsilon, v^*} \geq \text{esssup}_{u \in \mathcal{U}_{t, \tau^\delta}} Y_t^{2, u, v^*} - \varepsilon\delta.$$

From the above inequality and (4.7) it follows that

$$Y_t^{2, u^\varepsilon, v^*} \geq -C\delta^{\frac{3}{2}} - \varepsilon\delta. \quad (4.8)$$

Let us consider the following BSDEs:

$$\left\{ \begin{array}{l} Y_s^{3, u^\varepsilon, v^*} = \int_s^{t+\delta} F_j(r, x, Y_r^{3, u^\varepsilon, v^*}, Z_r^{3, u^\varepsilon, v^*}, u_r^\varepsilon, v_r^*) dr + K_{t+\delta}^{3, u^\varepsilon, v^*} - K_s^{3, u^\varepsilon, v^*} \\ \quad - \int_s^{t+\delta} Z_r^{3, u^\varepsilon, v^*} dB_r, \\ Y_s^{3, u^\varepsilon, v^*} \geq -\tau, \quad s \in [t, t + \delta], \\ \int_t^{t+\delta} (Y_r^{3, u^\varepsilon, v^*} + \tau) dr = 0, \quad K_t^{3, u^\varepsilon, v^*} = 0, \end{array} \right.$$

and

$$\left\{ \begin{array}{l} Y_s = \int_s^{t+\delta} (C|Y_r| + C|Z_r| - \theta)dr + K_{t+\delta} - K_s - \int_s^{t+\delta} Z_r dB_r, \\ Y_s \geq -\tau, \quad s \in [t, t+\delta], \quad K_t = 0 \\ \int_t^{t+\delta} (Y_s + \tau)ds = 0. \end{array} \right. \quad (4.9)$$

We choose δ such that $\delta \leq \delta'$ and $\frac{\theta}{C}(1 - \exp(-C\delta)) \leq \tau$. Then the solution of (4.9) is

$$Y_s = -\frac{\theta}{C}(1 - \exp(C(s - t - \delta))), \quad Z_s = 0, \quad K_s = 0, \quad s \in [t, t + \delta].$$

By Lemma 2.4 and (4.8) we see that

$$Y_t \geq Y_t^{3,u^\varepsilon,v^*} \geq Y_t^{2,u^\varepsilon,v^*}.$$

Consequently,

$$-\frac{\theta}{C}(1 - \exp(-C\delta)) \geq -C\delta^{\frac{3}{2}} - \varepsilon\delta,$$

i.e.,

$$-\frac{\theta}{C\delta}(1 - \exp(-C\delta)) \geq -C\delta^{\frac{1}{2}} - \varepsilon.$$

Finally, letting $\delta \rightarrow 0$ and $\varepsilon \rightarrow 0$ we conclude $\theta \leq 0$. This is a contradiction with $\theta > 0$. Consequently,

$$\inf_{v \in V} \sup_{u \in U} F(t, x, 0, 0, 0, u, v) \geq 0,$$

from which we conclude that W_j is a viscosity solution of (4.1). $\square$

Let us now give a comparison theorem for the viscosity solution of (4.1). We first introduce the following space:

$$\Theta : = \left\{ \varphi \in C([0, T] \times \mathbb{R}^n) : \text{there exists a constant } A > 0 \text{ such that} \right. \\ \left. \lim_{|x| \rightarrow \infty} |\varphi(t, x)| \exp\{-A[\log(|x|^2 + 1)^{\frac{1}{2}}]^2\} = 0, \text{ uniformly in } t \in [0, T] \right\}.$$

Theorem 4.7. *Under the assumptions (H3.1)–(H3.3), if an upper semicontinuous function $u_1 \in \Theta$ is a viscosity subsolution of (4.1) (resp., (4.2)), and a lower semicontinuous function $u_2 \in \Theta$ is a viscosity supersolution of (4.1) (resp., (4.2)), then we have the following:*

$$u_1(t, x) \leq u_2(t, x), \quad \text{for all } (t, x) \in [0, T] \times \mathbb{R}^n.$$

By means of the arguments in Barles, Buckdahn and Pardoux [17], we can give the proof of this theorem and the proof is omitted here.

Remark 4.8. *By Proposition 3.9 we see that W_j (resp., U_j) is a viscosity solution of linear growth. Therefore, from the above theorem we know that W_j (resp., U_j) is the unique viscosity solution in Θ of the system (4.1) (resp., (4.2)).*

Isaacs condition:

For all $(t, x, y, p, A, u, v) \in [0, T] \times \mathbb{R}^n \times \mathbb{R} \times \mathbb{R} \times \mathbb{R}^d \times \mathbb{S}^d \times U \times V$, $j = 1, 2$, we have

$$\begin{aligned} & \sup_{u \in U} \inf_{v \in V} \left\{ \frac{1}{2} \text{tr}(\sigma \sigma^T(t, x, u, v)A) + pb(t, x, u, v) + f_j(t, x, y, p\sigma(t, x, u, v), u, v) \right\} \\ &= \inf_{v \in V} \sup_{u \in U} \left\{ \frac{1}{2} \text{tr}(\sigma \sigma^T(t, x, u, v)A) + pb(t, x, u, v) + f_j(t, x, y, p\sigma(t, x, u, v), u, v) \right\}. \end{aligned} \quad (4.10)$$

Corollary 4.9. *Let Isaacs condition (4.10) hold. Then we have, for all $(t, x) \in [0, T] \times \mathbb{R}^n$,*

$$(U_1(t, x), U_2(t, x)) = (W_1(t, x), W_2(t, x)).$$

5 Nash equilibrium payoff

In this section we suppose that Isaacs condition (4.10) holds. In order to simplify arguments, let us also assume that the coefficients $b, \sigma, f_j, \Phi_j, f_j$ and h_j satisfy assumptions (H3.1)–(H3.4) and are bounded. By Isaacs condition (4.10) and Corollary 4.9 we see that, for $(t, x) \in [0, T] \times \mathbb{R}^n$,

$$W_1(t, x) = \operatorname{esssup}_{\alpha \in \mathcal{A}_{t,T}} \operatorname{essinf}_{\beta \in \mathcal{B}_{t,T}} J_1(t, x; \alpha, \beta) = \operatorname{essinf}_{\beta \in \mathcal{B}_{t,T}} \operatorname{esssup}_{\alpha \in \mathcal{A}_{t,T}} J_1(t, x; \alpha, \beta),$$

and

$$W_2(t, x) = \operatorname{essinf}_{\alpha \in \mathcal{A}_{t,T}} \operatorname{esssup}_{\beta \in \mathcal{B}_{t,T}} J_2(t, x; \alpha, \beta) = \operatorname{esssup}_{\beta \in \mathcal{B}_{t,T}} \operatorname{essinf}_{\alpha \in \mathcal{A}_{t,T}} J_2(t, x; \alpha, \beta). \quad (5.1)$$

Let us introduce the definition of the Nash equilibrium payoff of stochastic differential games (see Lin [8] or Buckdahn, Cardaliaguet and Rainer [7]).

Definition 5.1. A couple $(e_1, e_2) \in \mathbb{R}^2$ is called a Nash equilibrium payoff at the point (t, x) if for any $\varepsilon > 0$, there exists $(\alpha_\varepsilon, \beta_\varepsilon) \in \mathcal{A}_{t,T} \times \mathcal{B}_{t,T}$ such that, for all $(\alpha, \beta) \in \mathcal{A}_{t,T} \times \mathcal{B}_{t,T}$,

$$J_1(t, x; \alpha_\varepsilon, \beta_\varepsilon) \geq J_1(t, x; \alpha, \beta_\varepsilon) - \varepsilon, \quad J_2(t, x; \alpha_\varepsilon, \beta_\varepsilon) \geq J_2(t, x; \alpha_\varepsilon, \beta) - \varepsilon, \quad \mathbb{P} - a.s., \quad (5.2)$$

and

$$|\mathbb{E}[J_j(t, x; \alpha_\varepsilon, \beta_\varepsilon)] - e_j| \leq \varepsilon, \quad j = 1, 2.$$

From Lemma 3.5 it follows that the following lemma holds.

Lemma 5.2. For any $\varepsilon > 0$ and $(\alpha_\varepsilon, \beta_\varepsilon) \in \mathcal{A}_{t,T} \times \mathcal{B}_{t,T}$, (5.2) holds if and only if, for all $(u, v) \in \mathcal{U}_{t,T} \times \mathcal{V}_{t,T}$,

$$J_1(t, x; \alpha_\varepsilon, \beta_\varepsilon) \geq J_1(t, x; u, \beta_\varepsilon(u)) - \varepsilon, \quad J_2(t, x; \alpha_\varepsilon, \beta_\varepsilon) \geq J_2(t, x; \alpha_\varepsilon(v), v) - \varepsilon, \quad \mathbb{P} - a.s. \quad (5.3)$$

Before giving the main results in this sections we first introduce the following lemma.

Lemma 5.3. Let $(t, x) \in [0, T] \times \mathbb{R}^n$ and $u \in \mathcal{U}_{t,T}$ be arbitrarily fixed. Then,

(i) for all $\delta \in [0, T - t]$ and $\varepsilon > 0$, there exists an NAD strategy $\alpha \in \mathcal{A}_{t,T}$ such that, for all $v \in \mathcal{V}_{t,T}$,

$$\begin{aligned} \alpha(v) &= u, \text{ on } [t, t + \delta], \\ {}^2Y_{t+\delta}^{t,x;\alpha(v),v} &\leq W_2(t + \delta, X_{t+\delta}^{t,x;\alpha(v),v}) + \varepsilon, \quad \mathbb{P} - a.s. \end{aligned}$$

(ii) for all $\delta \in [0, T - t]$ and $\varepsilon > 0$, there exists an NAD strategy $\alpha \in \mathcal{A}_{t,T}$ such that, for all $v \in \mathcal{V}_{t,T}$,

$$\begin{aligned} \alpha(v) &= u, \text{ on } [t, t + \delta], \\ {}^1Y_{t+\delta}^{t,x;\alpha(v),v} &\geq W_1(t + \delta, X_{t+\delta}^{t,x;\alpha(v),v}) - \varepsilon, \quad \mathbb{P} - a.s. \end{aligned}$$

Using arguments similar to Lin [8] we can prove this lemma. The proof is omitted here. We also need the following lemma, which can be established by standard arguments for SDEs.

Lemma 5.4. There exists a positive constant C such that, for all $(u, v), (u', v') \in \mathcal{U}_{t,T} \times \mathcal{V}_{t,T}$, and for all $\mathcal{F}_\tau$ -stopping times $S : \Omega \rightarrow [t, T]$ with $X_S^{t,x;u,v} = X_S^{t,x;u',v'}$, $\mathbb{P} - a.s.$, it holds, for all real $\tau \in [t, T]$,

$$\mathbb{E} \left[\sup_{0 \leq s \leq \tau} |X_{(S+s) \wedge T}^{t,x;u,v} - X_{(S+s) \wedge T}^{t,x;u',v'}|^2 \middle| \mathcal{F}_t \right] \leq C\tau, \quad \mathbb{P} - a.s.$$

Let us now give one of main results in this section: the characterization theorem of Nash equilibrium payoffs for nonzero-sum stochastic differential games with reflection.

Theorem 5.5. *Let Isaacs condition (4.10) hold. Then, for $(t, x) \in [0, T] \times \mathbb{R}^n$, $(e_1, e_2) \in \mathbb{R}^2$ is a Nash equilibrium payoff at point (t, x) if and only if for all $\varepsilon > 0$, there exist $u^\varepsilon \in \mathcal{U}_{t,T}$ and $v^\varepsilon \in \mathcal{V}_{t,T}$ such that for all $s \in [t, T]$ and $j = 1, 2$,*

$$\mathbb{P}\left({}^jY_s^{t,x;u^\varepsilon,v^\varepsilon} \geq W_j(s, X_s^{t,x;u^\varepsilon,v^\varepsilon}) - \varepsilon \mid \mathcal{F}_t\right) \geq 1 - \varepsilon, \quad \mathbb{P} - a.s., \quad (5.4)$$

and

$$|\mathbb{E}[J_j(t, x; u^\varepsilon, v^\varepsilon)] - e_j| \leq \varepsilon. \quad (5.5)$$

Proof of Theorem 5.5: Sufficiency of (5.4) and (5.5).

Proof. For arbitrarily fixed $\varepsilon > 0$ and some $\varepsilon_0 > 0$ (ε_0 depends on ε and will be precise later), let us assume that $(u^{\varepsilon_0}, v^{\varepsilon_0}) \in \mathcal{U}_{t,T} \times \mathcal{V}_{t,T}$ satisfies (5.4) and (5.5), i.e., for all $s \in [t, T]$ and $j = 1, 2$,

$$\mathbb{P}\left({}^jY_s^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}} \geq W_j(s, X_s^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}}) - \varepsilon_0 \mid \mathcal{F}_t\right) \geq 1 - \varepsilon_0, \quad \mathbb{P} - a.s., \quad (5.6)$$

and

$$|\mathbb{E}[J_j(t, x; u^{\varepsilon_0}, v^{\varepsilon_0})] - e_j| \leq \varepsilon_0. \quad (5.7)$$

We fix some partition: $t = t_0 \leq t_1 \leq \dots \leq t_m = T$ of $[t, T]$ and put $\tau = \sup_i |t_i - t_{i+1}|$. Let us apply Lemma 5.3 to u^{ε_0} and $t + \delta = t_1, \dots, t_m$, successively. Then, for $\varepsilon_1 > 0$ (ε_1 depends on ε and is specified later) we have the existence of NAD strategies $\alpha_i \in \mathcal{A}_{t,T}$, $i = 1, \dots, m$, such that, for all $v \in \mathcal{V}_{t,T}$,

$$\begin{aligned} \alpha_i(v) &= u^{\varepsilon_0}, \text{ on } [t, t_i], \\ {}^2Y_{t_i}^{t,x;\alpha_i(v),v} &\leq W_2(t_i, X_{t_i}^{t,x;\alpha_i(v),v}) + \varepsilon_1, \quad \mathbb{P} - a.s. \end{aligned} \quad (5.8)$$

For all $v \in \mathcal{V}_{t,T}$, we set

$$\begin{aligned} S^v &= \inf \left\{ s \geq t \mid \lambda(\{r \in [t, s] : v_r \neq v_r^{\varepsilon_0}\}) > 0 \right\}, \\ t^v &= \inf \left\{ t_i \geq S^v \mid i = 1, \dots, m \right\} \wedge T. \end{aligned}$$

Here λ denotes the Lebesgue measure on the real line $\mathbb{R}$. We see that S^v and t^v are stopping times such that $S^v \leq t^v \leq S^v + \tau$.

We put

$$\alpha_\varepsilon(v) = \begin{cases} u^{\varepsilon_0}, & \text{on } [t, t^v], \\ \alpha_i(v), & \text{on } (t_i, T] \times \{t^v = t_i\}, 1 \leq i \leq m. \end{cases}$$

Then, α_ε is an NAD strategy. It follows from (5.8) that

$$\begin{aligned} {}^2Y_{t^v}^{t,x;\alpha_\varepsilon(v),v} &= \sum_{i=1}^m {}^2Y_{t_i}^{t,x;\alpha_\varepsilon(v),v} 1_{\{t^v=t_i\}} \\ &\leq \sum_{i=1}^m W_2(t_i, X_{t_i}^{t,x;\alpha_\varepsilon(v),v}) 1_{\{t^v=t_i\}} + \varepsilon_1 \\ &= W_2(t^v, X_{t^v}^{t,x;\alpha_\varepsilon(v),v}) + \varepsilon_1, \quad \mathbb{P} - a.s. \end{aligned} \quad (5.9)$$

Let us show that, for all $\varepsilon > 0$ and $v \in \mathcal{V}_{t,T}$,

$$J_2(t, x; \alpha_\varepsilon(v), v) \leq J_2(t, x; u^{\varepsilon_0}, v^{\varepsilon_0}) + \varepsilon, \quad \alpha_\varepsilon(v^{\varepsilon_0}) = u^{\varepsilon_0}. \quad (5.10)$$

Thanks to (5.9), from Lemmas 2.3 and 2.4 we see that there exists a positive constant C such that

$$\begin{aligned} J_2(t, x, \alpha_\varepsilon(v), v) &= {}^2G_{t,t^v}^{t,x;\alpha_\varepsilon(v),v} [2Y_{t^v}^{t,x,\alpha_\varepsilon(v),v}] \\ &\leq {}^2G_{t,t^v}^{t,x;\alpha_\varepsilon(v),v} [W_2(t^v, X_{t^v}^{t,x;\alpha_\varepsilon(v),v}) + \varepsilon_1] \\ &\leq {}^2G_{t,t^v}^{t,x;\alpha_\varepsilon(v),v} [W_2(t^v, X_{t^v}^{t,x;\alpha_\varepsilon(v),v})] + C\varepsilon_1. \end{aligned} \quad (5.11)$$

Therefore, from Lemma 2.3

$$\begin{aligned} &| {}^2G_{t,t^v}^{t,x;\alpha_\varepsilon(v),v} [W_2(t^v, X_{t^v}^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}})] - {}^2G_{t,t^v}^{t,x;\alpha_\varepsilon(v),v} [W_2(t^v, X_{t^v}^{t,x;\alpha_\varepsilon(v),v})] | \\ &\leq C\mathbb{E}[|W_2(t^v, X_{t^v}^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}}) - W_2(t^v, X_{t^v}^{t,x;\alpha_\varepsilon(v),v})|^2 | \mathcal{F}_t]^{\frac{1}{2}} \\ &\leq C\mathbb{E}[|X_{t^v}^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}} - X_{t^v}^{t,x;\alpha_\varepsilon(v),v}|^2 | \mathcal{F}_t]^{\frac{1}{2}} \\ &\leq C\tau^{\frac{1}{2}}, \quad \mathbb{P} - a.s. \end{aligned}$$

for the last two inequalities we have used Proposition 3.9 and Lemma 5.4. Then, (5.11) yield

$$\begin{aligned} &J_2(t, x, \alpha_\varepsilon(v), v) \\ &\leq {}^2G_{t,t^v}^{t,x;\alpha_\varepsilon(v),v} [W_2(t^v, X_{t^v}^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}})] + C\varepsilon_1 \\ &\quad + | {}^2G_{t,t^v}^{t,x;\alpha_\varepsilon(v),v} [W_2(t^v, X_{t^v}^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}})] - {}^2G_{t,t^v}^{t,x;\alpha_\varepsilon(v),v} [W_2(t^v, X_{t^v}^{t,x;\alpha_\varepsilon(v),v})] | \\ &\leq {}^2G_{t,t^v}^{t,x;\alpha_\varepsilon(v),v} [W_2(t^v, X_{t^v}^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}})] + C\varepsilon_1 + C\tau^{\frac{1}{2}}. \end{aligned}$$

Putting

$$\Omega_s = \left\{ {}^2Y_s^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}} \geq W_2(s, X_s^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}}) - \varepsilon_0 \right\}, s \in [t, T]. \quad (5.12)$$

we have

$$\begin{aligned} &J_2(t, x, \alpha_\varepsilon(v), v) \\ &\leq {}^2G_{t,t^v}^{t,x;\alpha_\varepsilon(v),v} \left[\sum_{i=1}^m W_2(t_i, X_{t_i}^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}}) 1_{\{t^v=t_i\}} \right] + C\varepsilon_1 + C\tau^{\frac{1}{2}} \\ &\leq {}^2G_{t,t^v}^{t,x;\alpha_\varepsilon(v),v} \left[\sum_{i=1}^m W_2(t_i, X_{t_i}^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}}) 1_{\{t^v=t_i\}} 1_{\Omega_{t_i}} \right] + C\varepsilon_1 + C\tau^{\frac{1}{2}} + I. \end{aligned} \quad (5.13)$$

where

$$I = | {}^2G_{t,t^v}^{t,x;\alpha_\varepsilon(v),v} \left[\sum_{i=1}^m W_2(t_i, X_{t_i}^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}}) 1_{\{t^v=t_i\}} \right] - {}^2G_{t,t^v}^{t,x;\alpha_\varepsilon(v),v} \left[\sum_{i=1}^m W_2(t_i, X_{t_i}^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}}) 1_{\{t^v=t_i\}} 1_{\Omega_{t_i}} \right] |$$

Since Φ_2, f_2 and h_2 are bounded, from Lemma 2.2 we conclude that W_2 is bounded. Therefore,

$$\begin{aligned} I &\leq \mathbb{E} \left[\sum_{i=1}^m |W_2(t_i, X_{t_i}^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}})|^2 1_{\{t^v=t_i\}} 1_{\Omega_{t_i}^c} \middle| \mathcal{F}_t \right]^{\frac{1}{2}} \\ &\leq C \sum_{i=1}^m \mathbb{P}(\Omega_{t_i}^c | \mathcal{F}_t)^{\frac{1}{2}} \leq Cm\varepsilon_0^{\frac{1}{2}}, \end{aligned} \quad (5.14)$$

where we have used (5.6) for the latter estimate. From the Lemmas 2.3, 2.4 and (5.12) we have

$$\begin{aligned} &{}^2G_{t,t^v}^{t,x;\alpha_\varepsilon(v),v} \left[\sum_{i=1}^m W_2(t_i, X_{t_i}^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}}) 1_{\{t^v=t_i\}} 1_{\Omega_{t_i}} \right] \\ &\leq {}^2G_{t,t^v}^{t,x;\alpha_\varepsilon(v),v} \left[\sum_{i=1}^m (2Y_{t_i}^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}} + \varepsilon_0) 1_{\{t^v=t_i\}} 1_{\Omega_{t_i}} \right] \end{aligned}$$

$$\begin{aligned}
&\leq {}^2G_{t,t^v}^{t,x;\alpha_\varepsilon(v),v} \left[\sum_{i=1}^m {}^2Y_{t_i}^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}} 1_{\{t^v=t_i\}} 1_{\Omega_{t_i}} + \varepsilon_0 \right] \\
&\leq {}^2G_{t,t^v}^{t,x;\alpha_\varepsilon(v),v} \left[\sum_{i=1}^m {}^2Y_{t_i}^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}} 1_{\{t^v=t_i\}} 1_{\Omega_{t_i}} \right] + C\varepsilon_0,
\end{aligned}$$

and using the above arguments we also have

$$\begin{aligned}
&| {}^2G_{t,t^v}^{t,x;\alpha_\varepsilon(v),v} \left[\sum_{i=1}^m {}^2Y_{t_i}^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}} 1_{\{t^v=t_i\}} 1_{\Omega_{t_i}} \right] - {}^2G_{t,t^v}^{t,x;\alpha_\varepsilon(v),v} \left[\sum_{i=1}^m {}^2Y_{t_i}^{t,x;\alpha_\varepsilon(v),v} 1_{\{t^v=t_i\}} \right] | \\
&\leq Cm\varepsilon_0^{\frac{1}{2}}.
\end{aligned}$$

Consequently,

$$\begin{aligned}
&{}^2G_{t,t^v}^{t,x;\alpha_\varepsilon(v),v} \left[\sum_{i=1}^m W_2(t_i, X_{t_i}^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}}) 1_{\{t^v=t_i\}} 1_{\Omega_{t_i}} \right] \\
&\leq {}^2G_{t,t^v}^{t,x;\alpha_\varepsilon(v),v} [{}^2Y_{t^v}^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}}] + C\varepsilon_0 + Cm\varepsilon_0^{\frac{1}{2}} \\
&\leq | {}^2G_{t,t^v}^{t,x;\alpha_\varepsilon(v),v} [{}^2Y_{t^v}^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}}] - {}^2G_{t,t^v}^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}} [{}^2Y_{t^v}^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}}] | \\
&\quad + {}^2G_{t,t^v}^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}} [{}^2Y_{t^v}^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}}] + C\varepsilon_0 + Cm\varepsilon_0^{\frac{1}{2}} \\
&= | {}^2G_{t,t^v}^{t,x;\alpha_\varepsilon(v),v} [{}^2Y_{t^v}^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}}] - {}^2G_{t,t^v}^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}} [{}^2Y_{t^v}^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}}] | \\
&\quad + J_2(t, x; u^{\varepsilon_0}, v^{\varepsilon_0}) + C\varepsilon_0 + Cm\varepsilon_0^{\frac{1}{2}} \\
&\leq J_2(t, x; u^{\varepsilon_0}, v^{\varepsilon_0}) + C\varepsilon_0 + Cm\varepsilon_0^{\frac{1}{2}} + C\tau^{\frac{1}{2}},
\end{aligned}$$

where we have used the fact that

$$| {}^2G_{t,t^v}^{t,x;\alpha_\varepsilon(v),v} [{}^2Y_{t^v}^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}}] - {}^2G_{t,t^v}^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}} [{}^2Y_{t^v}^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}}] | \leq C\tau^{\frac{1}{2}}.$$

Indeed, let us consider the following BSDE

$$\left\{ \begin{array}{l} y_s = {}^2Y_{t^v}^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}} + \int_s^{t^v} f_2(r, X_r^{t,x;\alpha_\varepsilon(v),v}, y_r, z_r, \alpha_\varepsilon(v)_r, v_r) dr + k_{t^v} - k_s - \int_s^{t^v} z_r dB_r, \\ y_s \geq h_2(s, X_s^{t,x;\alpha_\varepsilon(v),v}), \quad s \in [t, t^v], \\ k_t = 0, \quad \int_t^{t^v} (y_r - h_2(r, X_r^{t,x;\alpha_\varepsilon(v),v})) dk_r = 0, \end{array} \right.$$

which compared with

$$\left\{ \begin{array}{l} {}^2Y_s^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}} = {}^2Y_{t^v}^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}} + \int_s^{t^v} f_2(r, X_r^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}}, {}^2Y_r^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}}, {}^2Z_r^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}}, u_r^{\varepsilon_0}, v_r^{\varepsilon_0}) dr \\ \quad + {}^2K_{t^v}^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}} - {}^2K_s^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}} - \int_s^{t^v} {}^2Z_r^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}} dB_r, \\ {}^2Y_s^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}} \geq h_2(s, X_s^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}}), \quad s \in [t, t^v], \\ {}^2K_t^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}} = 0, \quad \int_t^{t^v} ({}^2Y_r^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}} - h_2(r, X_r^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}})) d {}^2K_r^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}} = 0, \end{array} \right.$$

Note that $\alpha_\varepsilon(v) = u^{\varepsilon_0}$, on $[[t, t^v]]$, $v = v^{\varepsilon_0}$, on $[[t, S^v]]$, and from Lemma 2.3 we obtain

$$\begin{aligned}
&| {}^2G_{t,t^v}^{t,x;\alpha_\varepsilon(v),v} [{}^2Y_{t^v}^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}}] - {}^2G_{t,t^v}^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}} [{}^2Y_{t^v}^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}}] |^2 \\
&\leq C\mathbb{E} \left[\int_t^{t^v} |f_2(r, X_r^{t,x;\alpha_\varepsilon(v),v}, y_r, z_r, \alpha_\varepsilon(v)_r, v_r) - f_2(r, X_r^{t,x;u^{\varepsilon_0},v^{\varepsilon_0}}, y_r, z_r, u_r^{\varepsilon_0}, v_r^{\varepsilon_0})|^2 | \mathcal{F}_t \right]
\end{aligned}$$

$$\begin{aligned}
& +C\mathbb{E}[\sup_{r \in [t, t^v]} |h_2(r, X_r^{t, x; \alpha_\varepsilon(v), v}) - h_2(r, X_r^{t, x; u^{\varepsilon_0}, v^{\varepsilon_0}})|^2 | \mathcal{F}_t]^\frac{1}{2} \\
= & C\mathbb{E}[\int_{S^v}^{t^v} |f_2(r, X_r^{t, x; \alpha_\varepsilon(v), v}, y_r, z_r, \alpha_\varepsilon(v)_r, v_r) - f_2(r, X_r^{t, x; u^{\varepsilon_0}, v^{\varepsilon_0}}, y_r, z_r, u_r^{\varepsilon_0}, v_r^{\varepsilon_0})|^2 | \mathcal{F}_t] \\
& +C\mathbb{E}[\sup_{r \in [S^v, t^v]} |X_r^{t, x; \alpha_\varepsilon(v), v} - X_r^{t, x; u^{\varepsilon_0}, v^{\varepsilon_0}}|^2 | \mathcal{F}_t]^\frac{1}{2} \\
\leq & C\mathbb{E}[\int_{S^v}^{t^v} 1_{\{v_r \neq v_r^{\varepsilon_0}\}} | \mathcal{F}_t] + C\tau^\frac{1}{2} \leq C\mathbb{E}[t^v - S^v | \mathcal{F}_t] + C\tau^\frac{1}{2} \leq C\tau^\frac{1}{2},
\end{aligned}$$

where we have used the boundedness of f_2, b and σ . Consequently,

$$\begin{aligned}
& {}^2G_{t, t^v}^{t, x; \alpha_\varepsilon(v), v} [\sum_{i=1}^m W_2(t_i, X_{t_i}^{t, x; u^{\varepsilon_0}, v^{\varepsilon_0}}) 1_{\{t^v = t_i\}} 1_{\Omega_{t_i}}] \\
& \leq C\tau^\frac{1}{2} + J_2(t, x; u^{\varepsilon_0}, v^{\varepsilon_0}) + C\varepsilon_0 + Cm\varepsilon_0^\frac{1}{2},
\end{aligned}$$

from where, (5.13) and (5.14) we see that

$$J_2(t, x; \alpha_\varepsilon(v), v) \leq J_2(t, x; u^{\varepsilon_0}, v^{\varepsilon_0}) + C\varepsilon_0 + Cm\varepsilon_0^\frac{1}{4} + C\varepsilon_1 + C\tau^\frac{1}{4}.$$

We can choose $\tau > 0, \varepsilon_0 > 0$, and $\varepsilon_1 > 0$ such that $C\varepsilon_0 + Cm\varepsilon_0^\frac{1}{2} + C\varepsilon_1 + C\tau^\frac{1}{4} \leq \varepsilon$ and $\varepsilon_0 < \varepsilon$. Thus,

$$J_2(t, x; \alpha_\varepsilon(v), v) \leq J_2(t, x; u^{\varepsilon_0}, v^{\varepsilon_0}) + \varepsilon, v \in \mathcal{V}_{t, T}.$$

Using a symmetric argument we can construct $\beta_\varepsilon \in \mathcal{B}_{t, T}$ such that, for all $u \in \mathcal{U}_{t, T}$,

$$J_1(t, x; u, \beta_\varepsilon(u)) \leq J_1(t, x; u^{\varepsilon_0}, v^{\varepsilon_0}) + \varepsilon, \quad \beta_\varepsilon(u^{\varepsilon_0}) = v^{\varepsilon_0}. \quad (5.15)$$

Finally, from (5.10), (5.15), (5.7) and Lemma 5.2 it follows that $(\alpha_\varepsilon, \beta_\varepsilon)$ satisfies Definition 5.1. Hence, (e_1, e_2) is a Nash equilibrium payoff. $\square$

Proof of Theorem 5.5: Necessity of (5.4) and (5.5).

Proof. Let us suppose that $(e_1, e_2) \in \mathbb{R}^2$ is a Nash equilibrium payoff at the point (t, x) . Then, for any sufficiently small $\varepsilon > 0$, there exists $(\alpha_\varepsilon, \beta_\varepsilon) \in \mathcal{A}_{t, T} \times \mathcal{B}_{t, T}$ such that, for all $(\alpha, \beta) \in \mathcal{A}_{t, T} \times \mathcal{B}_{t, T}$

$$J_1(t, x; \alpha_\varepsilon, \beta_\varepsilon) \geq J_1(t, x; \alpha, \beta_\varepsilon) - \varepsilon^4, \quad J_2(t, x; \alpha_\varepsilon, \beta_\varepsilon) \geq J_2(t, x; \alpha_\varepsilon, \beta) - \varepsilon^4, \quad \mathbb{P} - a.s., \quad (5.16)$$

and

$$|\mathbb{E}[J_j(t, x; \alpha_\varepsilon, \beta_\varepsilon)] - e_j| \leq \varepsilon^4, \quad j = 1, 2. \quad (5.17)$$

It follows from Lemma 3.5 that there exists a unique couple $(u^\varepsilon, v^\varepsilon)$ such that

$$\alpha_\varepsilon(v^\varepsilon) = u^\varepsilon, \beta_\varepsilon(u^\varepsilon) = v^\varepsilon.$$

We give the proof by contradiction. Note that (5.17) means that (5.5) holds. Assuming that (5.4) doesn't hold true, for all $\varepsilon' > 0$, there exists some $\varepsilon \in (0, \varepsilon')$ (for which we use the notations introduced above) and $\delta \in [0, T - t]$ such that, for some $j \in \{1, 2\}$, say for $j = 1$,

$$\mathbb{P}(\mathbb{P}(^1Y_{t+\delta}^{t, x; u^\varepsilon, v^\varepsilon} < W_1(s, X_{t+\delta}^{t, x; u^\varepsilon, v^\varepsilon}) - \varepsilon \mid \mathcal{F}_t) > \varepsilon) > 0. \quad (5.18)$$

We apply Lemma 5.3 to u^ε and $t + \delta$, it follows that, there exists an NAD strategy $\tilde{\alpha} \in \mathcal{A}_{t, T}$, such that, for all $v \in \mathcal{V}_{t, T}$,

$$\tilde{\alpha}(v) = u^\varepsilon, \text{ on } [t, t + \delta],$$

$${}^1Y_{t+\delta}^{t,x;\tilde{\alpha}(v),v} \geq W_1(t+\delta, X_{t+\delta}^{t,x;\tilde{\alpha}(v),v}) - \frac{\varepsilon}{2}, \quad \mathbb{P} - a.s. \quad (5.19)$$

It follows from Lemma 3.5 that there exists a unique couple $(u, v) \in \mathcal{U}_{t,T} \times \mathcal{V}_{t,T}$ such that

$$\tilde{\alpha}(v) = u, \quad \beta_\varepsilon(u) = v.$$

Let us

$$A = \left\{ {}^1Y_{t+\delta}^{t,x;u^\varepsilon,v^\varepsilon} < W_1(t+\delta, X_{t+\delta}^{t,x;u^\varepsilon,v^\varepsilon}) - \varepsilon \right\} \in \mathcal{F}_{t+\delta}, \quad (5.20)$$

and define

$$\tilde{u} = \begin{cases} u^\varepsilon, & \text{on } ([t, t+\delta) \times \Omega) \cup ([t+\delta, T] \times A^c), \\ u, & \text{on } [t+\delta, T] \times A. \end{cases}$$

Thanks to the nonanticipativity of $\beta_\varepsilon \in \mathcal{B}_{t,T}$ we see that $\beta_\varepsilon(\tilde{u}) = \beta_\varepsilon(u^\varepsilon) = v^\varepsilon$ on $[t, t+\delta]$, and for all $s \in [t+\delta, T]$,

$$\beta_\varepsilon(\tilde{u}) = \begin{cases} \beta_\varepsilon(u) = v, & \text{on } A, \\ \beta_\varepsilon(u^\varepsilon) = v^\varepsilon, & \text{on } A^c. \end{cases}$$

From standard arguments for SDEs and BSDEs it follows that

$$X^{t,x;\tilde{u},\beta_\varepsilon(\tilde{u})} = X^{t,x;u^\varepsilon,v^\varepsilon}, \quad \text{on } [t, t+\delta],$$

$$X^{t,x;\tilde{u},\beta_\varepsilon(\tilde{u})} = \begin{cases} X^{t,x;\tilde{\alpha}(v),v}, & \text{on } [t+\delta, T] \times A, \\ X^{t,x;u^\varepsilon,v^\varepsilon}, & \text{on } [t+\delta, T] \times A^c, \end{cases}$$

as well as

$${}^1Y_{t+\delta}^{t,x;\tilde{u},\beta_\varepsilon(\tilde{u})} = \begin{cases} {}^1Y_{t+\delta}^{t,x;\tilde{\alpha}(v),v}, & \text{on } A, \\ {}^1Y_{t+\delta}^{t,x;u^\varepsilon,v^\varepsilon}, & \text{on } A^c. \end{cases}$$

Consequently,

$$\begin{aligned} J_1(t, x; \tilde{u}, \beta_\varepsilon(\tilde{u})) &= {}^1Y_t^{t,x;\tilde{u},\beta_\varepsilon(\tilde{u})} = {}^1G_{t,t+\delta}^{t,x;\tilde{u},\beta_\varepsilon(\tilde{u})} [{}^1Y_{t+\delta}^{t,x;\tilde{u},\beta_\varepsilon(\tilde{u})}] \\ &= {}^1G_{t,t+\delta}^{t,x;\tilde{u},\beta_\varepsilon(\tilde{u})} [{}^1Y_{t+\delta}^{t,x;\tilde{u},\beta_\varepsilon(\tilde{u})} 1_A + {}^1Y_{t+\delta}^{t,x;\tilde{u},\beta_\varepsilon(\tilde{u})} 1_{A^c}] \\ &= {}^1G_{t,t+\delta}^{t,x;\tilde{u},\beta_\varepsilon(\tilde{u})} [{}^1Y_{t+\delta}^{t,x;\tilde{\alpha}(v),v} 1_A + {}^1Y_{t+\delta}^{t,x;u^\varepsilon,v^\varepsilon} 1_{A^c}]. \end{aligned}$$

By Lemma 2.4 and (5.19) we deduce that

$$\begin{aligned} J_1(t, x; \tilde{u}, \beta_\varepsilon(\tilde{u})) &= {}^1G_{t,t+\delta}^{t,x;\tilde{u},\beta_\varepsilon(\tilde{u})} [{}^1Y_{t+\delta}^{t,x;\tilde{\alpha}(v),v} 1_A + {}^1Y_{t+\delta}^{t,x;u^\varepsilon,v^\varepsilon} 1_{A^c}] \\ &\geq {}^1G_{t,t+\delta}^{t,x;\tilde{u},\beta_\varepsilon(\tilde{u})} [(W_1(t+\delta, X_{t+\delta}^{t,x;\tilde{\alpha}(v),v}) - \frac{\varepsilon}{2}) 1_A + {}^1Y_{t+\delta}^{t,x;u^\varepsilon,v^\varepsilon} 1_{A^c}] \\ &= {}^1G_{t,t+\delta}^{t,x;\tilde{u},\beta_\varepsilon(\tilde{u})} [W_1(t+\delta, X_{t+\delta}^{t,x;u^\varepsilon,v^\varepsilon}) 1_A + {}^1Y_{t+\delta}^{t,x;u^\varepsilon,v^\varepsilon} 1_{A^c} - \frac{\varepsilon}{2} 1_A]. \end{aligned}$$

Therefore, from (5.20)

$$\begin{aligned} J_1(t, x; \tilde{u}, \beta_\varepsilon(\tilde{u})) &\geq {}^1G_{t,t+\delta}^{t,x;\tilde{u},\beta_\varepsilon(\tilde{u})} [W_1(t+\delta, X_{t+\delta}^{t,x;u^\varepsilon,v^\varepsilon}) 1_A + {}^1Y_{t+\delta}^{t,x;u^\varepsilon,v^\varepsilon} 1_{A^c} - \frac{\varepsilon}{2} 1_A] \\ &\geq {}^1G_{t,t+\delta}^{t,x;\tilde{u},\beta_\varepsilon(\tilde{u})} [({}^1Y_{t+\delta}^{t,x;u^\varepsilon,v^\varepsilon} + \varepsilon) 1_A + {}^1Y_{t+\delta}^{t,x;u^\varepsilon,v^\varepsilon} 1_{A^c} - \frac{\varepsilon}{2} 1_A] \\ &= {}^1G_{t,t+\delta}^{t,x;\tilde{u},\beta_\varepsilon(\tilde{u})} [{}^1Y_{t+\delta}^{t,x;u^\varepsilon,v^\varepsilon} + \frac{\varepsilon}{2} 1_A] \end{aligned}$$

$$= {}^1G_{t,t+\delta}^{t,x;u^\varepsilon,v^\varepsilon} [{}^1Y_{t+\delta}^{t,x;u^\varepsilon,v^\varepsilon} + \frac{\varepsilon}{2}1_A]. \quad (5.21)$$

Let us consider the following BSDE:

$$\left\{ \begin{array}{l} y_s = {}^1Y_{t+\delta}^{t,x;u^\varepsilon,v^\varepsilon} + \frac{\varepsilon}{2}1_A + \int_s^{t+\delta} f_1(r, X_r^{t,x;u^\varepsilon,v^\varepsilon}, y_r, z_r, u_r^\varepsilon, v_r^\varepsilon) dr + k_{t+\delta} - k_s - \int_s^{t+\delta} z_r dB_r, \\ y_s \geq h_1(s, X_s^{t,x;u^\varepsilon,v^\varepsilon}), \quad s \in [t, t+\delta], \\ k_t = 0, \quad \int_t^{t+\delta} (y_s - h_1(s, X_s^{t,x;u^\varepsilon,v^\varepsilon})) dk_r = 0, \end{array} \right.$$

as well as

$$\left\{ \begin{array}{l} {}^1Y_s^{t,x;u^\varepsilon,v^\varepsilon} = {}^1Y_{t+\delta}^{t,x;u^\varepsilon,v^\varepsilon} + \int_s^{t+\delta} f_1(r, X_r^{t,x;u^\varepsilon,v^\varepsilon}, {}^1Y_r^{t,x;u^\varepsilon,v^\varepsilon}, {}^1Z_r^{t,x;u^\varepsilon,v^\varepsilon}, u_r^\varepsilon, v_r^\varepsilon) dr \\ \quad + {}^1K_{t+\delta}^{t,x;u^\varepsilon,v^\varepsilon} - {}^1K_s^{t,x;u^\varepsilon,v^\varepsilon} - \int_s^{t+\delta} {}^1Z_r^{t,x;u^\varepsilon,v^\varepsilon} dB_r, \\ {}^1Y_s^{t,x;u^\varepsilon,v^\varepsilon} \geq h_1(s, X_s^{t,x;u^\varepsilon,v^\varepsilon}), \quad s \in [t, t+\delta], \\ {}^1K_t^{t,x;u^\varepsilon,v^\varepsilon} = 0, \quad \int_t^{t+\delta} ({}^1Y_r^{t,x;u^\varepsilon,v^\varepsilon} - h_1(r, X_r^{t,x;u^\varepsilon,v^\varepsilon})) d {}^1K_r^{t,x;u^\varepsilon,v^\varepsilon} = 0, \end{array} \right.$$

Let us set

$$\bar{y}_s = y_s - {}^1Y_s^{t,x;u^\varepsilon,v^\varepsilon}, \quad \bar{k}_s = k_s - {}^1K_s^{t,x;u^\varepsilon,v^\varepsilon}, \quad \bar{z}_s = z_s - {}^1Z_s^{t,x;u^\varepsilon,v^\varepsilon}, \quad s \in [t, t+\delta].$$

Then from Lemma 2.4 it follows that $\bar{k}_s \geq 0$, and we have

$$\begin{aligned} \bar{y}_s &= \int_s^{t+\delta} [f_1(r, X_r^{t,x;u^\varepsilon,v^\varepsilon}, y_r, z_r, u_r^\varepsilon, v_r^\varepsilon) - f_1(r, X_r^{t,x;u^\varepsilon,v^\varepsilon}, {}^1Y_r^{t,x;u^\varepsilon,v^\varepsilon}, {}^1Z_r^{t,x;u^\varepsilon,v^\varepsilon}, u_r^\varepsilon, v_r^\varepsilon)] dr \\ &\quad + \bar{k}_{t+\delta} - \bar{k}_s + \frac{\varepsilon}{2}1_A - \int_s^{t+\delta} \bar{z}_r dB_r, \quad s \in [t, t+\delta]. \end{aligned} \quad (5.22)$$

For $r \in [t, t+\delta]$, we set

$$\begin{aligned} a_r &= 1_{\{\bar{y}_r \neq 0\}} (\bar{y}_r)^{-1} \left(f_1(r, X_r^{t,x;u^\varepsilon,v^\varepsilon}, y_r, z_r, u_r^\varepsilon, v_r^\varepsilon) - f_1(r, X_r^{t,x;u^\varepsilon,v^\varepsilon}, {}^1Y_r^{t,x;u^\varepsilon,v^\varepsilon}, {}^1Z_r^{t,x;u^\varepsilon,v^\varepsilon}, u_r^\varepsilon, v_r^\varepsilon) \right), \\ b_r &= 1_{\{\bar{z}_r \neq 0\}} (\bar{z}_r)^{-1} \left(f_1(r, X_r^{t,x;u^\varepsilon,v^\varepsilon}, {}^1Y_r^{t,x;u^\varepsilon,v^\varepsilon}, {}^1Z_r^{t,x;u^\varepsilon,v^\varepsilon}, u_r^\varepsilon, v_r^\varepsilon) \right. \\ &\quad \left. - f_1(r, X_r^{t,x;u^\varepsilon,v^\varepsilon}, {}^1Y_r^{t,x;u^\varepsilon,v^\varepsilon}, {}^1Z_r^{t,x;u^\varepsilon,v^\varepsilon}, u_r^\varepsilon, v_r^\varepsilon) \right). \end{aligned}$$

Then, $|a_r| \leq L$, $|b_r| \leq L$, $r \in [t, t+\delta]$, and BSDE (5.22) has the following form:

$$\bar{y}_s = \frac{\varepsilon}{2}1_A + \int_s^{t+\delta} [a_r \bar{y}_r + b_r \bar{z}_r] dr - \int_s^{t+\delta} \bar{z}_r dB_r + \bar{k}_{t+\delta} - \bar{k}_s, \quad s \in [t, t+\delta].$$

Put

$$Q_s = \exp \left(\int_t^s a_r dr + \int_t^s b_r dB_r - \frac{1}{2} \int_t^s |b_r|^2 dr \right), \quad s \in [t, t+\delta].$$

It follows from Itô's formula to $\bar{y}_s Q_s$ that

$$\bar{y}_t = \frac{\varepsilon}{2} \mathbb{E}[1_A Q_{t+\delta} + \int_t^{t+\delta} Q_r d\bar{k}_r \mid \mathcal{F}_t] \geq \frac{\varepsilon}{2} \mathbb{E}[1_A Q_{t+\delta} \mid \mathcal{F}_t].$$

Thanks to the Hölder inequality we obtain

$$\mathbb{P}(A \mid \mathcal{F}_t)^2 = (\mathbb{E}[1_A \mid \mathcal{F}_t])^2 \leq \mathbb{E}[1_A Q_{t+\delta} \mid \mathcal{F}_t] \mathbb{E}[Q_{t+\delta}^{-1} \mid \mathcal{F}_t].$$

We note that

$$\begin{aligned}
\mathbb{E}[Q_{t+\delta}^{-1} | \mathcal{F}_t] &= \mathbb{E}[\exp\left(-\int_t^{t+\delta} a_r dr - \int_t^{t+\delta} b_r dB_r + \frac{1}{2} \int_t^{t+\delta} |b_r|^2 dr\right) | \mathcal{F}_t] \\
&\leq \exp(L\delta + L^2\delta) \mathbb{E}[\exp\left(-\int_t^{t+\delta} b_r dB_r - \frac{1}{2} \int_t^{t+\delta} |b_r|^2 dr\right) | \mathcal{F}_t] \\
&= \exp(L\delta + L^2\delta).
\end{aligned}$$

Let

$$\Delta = \left\{ \mathbb{P}({}^1Y_{t+\delta}^{t,x;u^\varepsilon,v^\varepsilon} < W_1(s, X_{t+\delta}^{t,x;u^\varepsilon,v^\varepsilon}) - \varepsilon \mid \mathcal{F}_t) > \varepsilon \right\} \left(= \{ \mathbb{P}(A | \mathcal{F}_t) > \varepsilon \} \right).$$

Then,

$$\begin{aligned}
\bar{y}_t &\geq \frac{\varepsilon}{2} \mathbb{E}[1_A Q_{t+\delta} | \mathcal{F}_t] \\
&\geq \frac{\exp(-L\delta - L^2\delta) \varepsilon}{2} (\mathbb{E}[1_A | \mathcal{F}_t])^2 = \frac{\exp(-L\delta - L^2\delta) \varepsilon}{2} (\mathbb{P}(A | \mathcal{F}_t))^2 \\
&> \frac{\varepsilon^3}{2} C_0 1_\Delta,
\end{aligned} \tag{5.23}$$

for $C_0 = \exp(-L\delta - L^2\delta)$, where we use (5.18) in the last inequality. Combining (5.23) with

$$\bar{y}_t = y_t - {}^1Y_t^{t,x;u^\varepsilon,v^\varepsilon} = {}^1G_{t,t+\delta}^{t,x;u^\varepsilon,v^\varepsilon} [{}^1Y_{t+\delta}^{t,x;u^\varepsilon,v^\varepsilon} + \frac{\varepsilon}{2} 1_A] - {}^1G_{t,t+\delta}^{t,x;u^\varepsilon,v^\varepsilon} [{}^1Y_{t+\delta}^{t,x;u^\varepsilon,v^\varepsilon}],$$

we have

$${}^1G_{t,t+\delta}^{t,x;u^\varepsilon,v^\varepsilon} [{}^1Y_{t+\delta}^{t,x;u^\varepsilon,v^\varepsilon} + \frac{\varepsilon}{2} 1_A] > {}^1G_{t,t+\delta}^{t,x;u^\varepsilon,v^\varepsilon} [{}^1Y_{t+\delta}^{t,x;u^\varepsilon,v^\varepsilon}] + \frac{\varepsilon^3}{2} C_0 1_\Delta,$$

and (5.21) then yields

$$J_1(t, x; \tilde{u}, \beta_\varepsilon(\tilde{u})) > J_1(t, x; \alpha_\varepsilon, \beta_\varepsilon) + \frac{\varepsilon^3}{2} C_0 1_\Delta.$$

Let us choose ε' sufficiently small such that $\frac{\varepsilon^3}{2} C_0 > \varepsilon^4$. Since $\mathbb{P}(\Delta) > 0$, the above inequality contradicts with (5.16) for $\alpha(\cdot) = \tilde{u}$. The proof is complete. $\square$

Before giving the existence theorem of a Nash equilibrium payoff we first establish the following proposition, which is crucial for the proof of the existence theorem of a Nash equilibrium payoff.

Proposition 5.6. *Under the assumptions of Theorem 5.5, for all $\varepsilon > 0$, there exists $(u^\varepsilon, v^\varepsilon) \in \mathcal{U}_{t,T} \times \mathcal{V}_{t,T}$ independent of $\mathcal{F}_t$ such that, for all $t \leq s_1 \leq s_2 \leq T$, $j = 1, 2$,*

$$\mathbb{P}\left(W_j(s_1, X_{s_1}^{t,x;u^\varepsilon,v^\varepsilon}) - \varepsilon \leq {}^jG_{s_1,s_2}^{t,x;u^\varepsilon,v^\varepsilon} [W_j(s_2, X_{s_2}^{t,x;u^\varepsilon,v^\varepsilon})] \mid \mathcal{F}_t \right) > 1 - \varepsilon.$$

Let us first give some preliminary result. Since the proof of the following lemma is similar to that in [8], we omit here.

Lemma 5.7. *For all $\varepsilon > 0$, all $\delta \in [0, T - t]$ and $x \in \mathbb{R}^n$, there exists $(u^\varepsilon, v^\varepsilon) \in \mathcal{U}_{t,T} \times \mathcal{V}_{t,T}$ independent of $\mathcal{F}_t$, such that, $j = 1, 2$,*

$$W_j(t, x) - \varepsilon \leq {}^jG_{t,t+\delta}^{t,x;u^\varepsilon,v^\varepsilon} [W_j(t + \delta, X_{t+\delta}^{t,x;u^\varepsilon,v^\varepsilon})], \quad \mathbb{P} - a.s.$$

Let us establish the following Lemma.

Lemma 5.8. *Let $n \geq 1$ and let us fix some partition $t = t_0 < t_1 < \dots < t_n = T$ of the interval $[t, T]$. Then, for all $\varepsilon > 0$, there exists $(u^\varepsilon, v^\varepsilon) \in \mathcal{U}_{t,T} \times \mathcal{V}_{t,T}$ independent of $\mathcal{F}_t$, such that, for all $i = 0, \dots, n-1$,*

$$W_j(t_i, X_{t_i}^{t,x;u^\varepsilon,v^\varepsilon}) - \varepsilon \leq {}^jG_{t_i,t_{i+1}}^{t,x;u^\varepsilon,v^\varepsilon} [W_j(t_{i+1}, X_{t_{i+1}}^{t,x;u^\varepsilon,v^\varepsilon})], \quad \mathbb{P} - a.s.$$

Proof. Let us prove this lemma by induction. By the above lemma, it is obvious for $i = 0$. We now suppose that $(u^\varepsilon, v^\varepsilon)$ independent of $\mathcal{F}_t$, is constructed on the interval $[t, t_i)$ and we shall define it on $[t_i, t_{i+1})$. From the above lemma it follows that, for all $y \in \mathbb{R}^n$, there exists $(u^y, v^y) \in \mathcal{U}_{t_i, T} \times \mathcal{V}_{t_i, T}$ independent of $\mathcal{F}_t$, such that,

$$W_j(t_i, y) - \frac{\varepsilon}{2} \leq {}^j G_{t_i, t_{i+1}}^{t_i, y; u^y, v^y} [W_j(t_{i+1}, X_{t_{i+1}}^{t_i, y; u^y, v^y})], \quad \mathbb{P} - a.s, j = 1, 2. \quad (5.24)$$

For arbitrarily $j = 1, 2$, for all $y, z \in \mathbb{R}^n$ and $s \in [t_i, t_{i+1}]$, we set

$$y_s^1 = {}^j G_{s, t_{i+1}}^{t_i, y; u^y, v^y} [W_j(t_{i+1}, X_{t_{i+1}}^{t_i, y; u^y, v^y})], \text{ and } y_s^2 = {}^j G_{s, t_{i+1}}^{t_i, z; u^y, v^y} [W_j(t_{i+1}, X_{t_{i+1}}^{t_i, z; u^y, v^y})].$$

Then let us consider the following BSDEs:

$$\left\{ \begin{array}{l} y_s^1 = W_j(t_{i+1}, X_{t_{i+1}}^{t_i, y; u^y, v^y}) + \int_s^{t_{i+1}} f_j(r, X_r^{t_i, y; u^y, v^y}, y_r^1, z_r^1, u_r^y, v_r^y) dr \\ \quad + {}^1 K_{t+\delta} - {}^1 K_s - \int_s^{t_{i+1}} z_r^1 dB_r, \\ y_s^1 \geq h_j(s, X_s^{t_i, y; u^y, v^y}), \quad s \in [t_i, t_{i+1}], \\ {}^1 K_{t_i} = 0, \quad \int_{t_i}^{t_{i+1}} (y_s^1 - h_j(s, X_s^{t_i, y; u^y, v^y})) d {}^1 K_r = 0, \end{array} \right.$$

and

$$\left\{ \begin{array}{l} y_s^1 = W_j(t_{i+1}, X_{t_{i+1}}^{t_i, z; u^y, v^y}) + \int_s^{t_{i+1}} f_j(r, X_r^{t_i, z; u^y, v^y}, y_r^1, z_r^1, u_r^y, v_r^y) dr \\ \quad + {}^2 K_{t+\delta} - {}^2 K_s - \int_s^{t_{i+1}} z_r^2 dB_r, \\ y_s^2 \geq h_j(s, X_s^{t_i, z; u^y, v^y}), \quad s \in [t_i, t_{i+1}], \\ {}^2 K_{t_i} = 0, \quad \int_{t_i}^{t_{i+1}} (y_s^2 - h_j(s, X_s^{t_i, z; u^y, v^y})) d {}^2 K_r = 0, \end{array} \right.$$

From the Lemmas 2.3 and 3.1 it follows that

$$\begin{aligned} & |{}^j G_{t_i, t_{i+1}}^{t_i, y; u^y, v^y} [W_j(t_{i+1}, X_{t_{i+1}}^{t_i, y; u^y, v^y})] - {}^j G_{t_i, t_{i+1}}^{t_i, z; u^y, v^y} [W_j(t_{i+1}, X_{t_{i+1}}^{t_i, z; u^y, v^y})]|^2 \\ & \leq C \mathbb{E}[|W_j(t_{i+1}, X_{t_{i+1}}^{t_i, y; u^y, v^y}) - W_j(t_{i+1}, X_{t_{i+1}}^{t_i, z; u^y, v^y})|^2 | \mathcal{F}_{t_i}] \\ & \quad + C \mathbb{E}[\left| \int_{t_i}^{t_{i+1}} f_j(r, X_r^{t_i, y; u^y, v^y}, y_r^1, z_r^1, u_r^y, v_r^y) dr - \int_{t_i}^{t_{i+1}} f_j(r, X_r^{t_i, z; u^y, v^y}, y_r^1, z_r^1, u_r^y, v_r^y) dr \right|^2 | \mathcal{F}_{t_i}] \\ & \quad + C \mathbb{E}[\sup_{t_i \leq s \leq t_{i+1}} |h_j(s, X_s^{t_i, y; u^y, v^y}) - h_j(s, X_s^{t_i, z; u^y, v^y})|^2 | \mathcal{F}_{t_i}]^{\frac{1}{2}} \\ & \leq C \mathbb{E}[|X_{t_{i+1}}^{t_i, y; u^y, v^y} - X_{t_{i+1}}^{t_i, z; u^y, v^y}|^2 | \mathcal{F}_t] + C \mathbb{E}[\left| \int_{t_i}^{t_{i+1}} |X_r^{t_i, y; u^y, v^y} - X_r^{t_i, z; u^y, v^y}|^2 dr \right| \mathcal{F}_{t_i}] \\ & \quad + C \mathbb{E}[\sup_{t_i \leq s \leq t_{i+1}} |X_s^{t_i, y; u^y, v^y} - X_s^{t_i, z; u^y, v^y}|^2 | \mathcal{F}_{t_i}]^{\frac{1}{2}} \\ & \leq C |y - z|. \end{aligned}$$

Combining the above inequality, Proposition 3.9 and (5.24) we see that

$$\begin{aligned} W_j(t_i, z) - \varepsilon & \leq W_j(t_i, y) - \varepsilon + C |y - z|^{\frac{1}{2}} \\ & \leq {}^j G_{t_i, t_{i+1}}^{t_i, y; u^y, v^y} [W_j(t_{i+1}, X_{t_{i+1}}^{t_i, y; u^y, v^y})] - \frac{\varepsilon}{2} + C |y - z|^{\frac{1}{2}} \\ & \leq {}^j G_{t_i, t_{i+1}}^{t_i, z; u^y, v^y} [W_j(t_{i+1}, X_{t_{i+1}}^{t_i, z; u^y, v^y})] - \frac{\varepsilon}{2} + C |y - z|^{\frac{1}{2}} \end{aligned}$$

$$\leq {}^j G_{t_i, t_{i+1}}^{t, z; u^y, v^y} [W_j(t_{i+1}, X_{t_{i+1}}^{t, z; u^y, v^y})], \mathbb{P} - a.s.,$$

for $C|y - z|^{\frac{1}{2}} \leq \frac{\varepsilon}{2}$.

Let $\{O_i\}_{i \geq 1} \subset \mathcal{B}(\mathbb{R}^n)$ be a partition of $\mathbb{R}^n$ with $\text{diam}(O_i) < \frac{\varepsilon}{2C}$ and let $y_l \in O_l$. Then, for $z \in O_l$,

$$W_j(t_i, z) - \varepsilon \leq {}^j G_{t_i, t_{i+1}}^{t, z; u^{y_l}, v^{y_l}} [W_j(t_{i+1}, X_{t_{i+1}}^{t, z; u^{y_l}, v^{y_l}})], \mathbb{P} - a.s. \quad (5.25)$$

Let us put

$$u^\varepsilon = \sum_{l \geq 1} 1_{O_l}(X^{t, x; u^\varepsilon, v^\varepsilon}) u^{y_l}, \quad v^\varepsilon = \sum_{l \geq 1} 1_{O_l}(X^{t, x; u^\varepsilon, v^\varepsilon}) v^{y_l}.$$

Then

$$\begin{aligned} & {}^j G_{t_i, t_{i+1}}^{t, x; u^\varepsilon, v^\varepsilon} [W_j(t_{i+1}, X_{t_{i+1}}^{t, x; u^\varepsilon, v^\varepsilon})] \\ &= {}^j G_{t_i, t_{i+1}}^{t_i, X_{t_i}^{t, x; u^\varepsilon, v^\varepsilon}; u^\varepsilon, v^\varepsilon} \left[\sum_{l \geq 1} W_j(t_{i+1}, X_{t_{i+1}}^{t_i, X_{t_i}^{t, x; u^\varepsilon, v^\varepsilon}; u^\varepsilon, v^\varepsilon}) 1_{O_l}(X_{t_i}^{t, x; u^\varepsilon, v^\varepsilon}) \right] \\ &= {}^j G_{t_i, t_{i+1}}^{t_i, X_{t_i}^{t, x; u^\varepsilon, v^\varepsilon}; u^\varepsilon, v^\varepsilon} \left[\sum_{l \geq 1} W_j(t_{i+1}, X_{t_{i+1}}^{t_i, X_{t_i}^{t, x; u^\varepsilon, v^\varepsilon}; u^{y_l}, v^{y_l}) 1_{O_l}(X_{t_i}^{t, x; u^\varepsilon, v^\varepsilon}) \right] \\ &= \sum_{l \geq 1} {}^j G_{t_i, t_{i+1}}^{t_i, X_{t_i}^{t, x; u^\varepsilon, v^\varepsilon}; u^{y_l}, v^{y_l}} [W_j(t_{i+1}, X_{t_{i+1}}^{t_i, X_{t_i}^{t, x; u^\varepsilon, v^\varepsilon}; u^{y_l}, v^{y_l})] 1_{O_l}(X_{t_i}^{t, x; u^\varepsilon, v^\varepsilon}), \end{aligned}$$

which together with (5.25) yields

$$\begin{aligned} & {}^j G_{t_i, t_{i+1}}^{t, x; u^\varepsilon, v^\varepsilon} [W_j(t_{i+1}, X_{t_{i+1}}^{t, x; u^\varepsilon, v^\varepsilon})] \\ &\geq \sum_{l \geq 1} [W_j(t_i, X_{t_i}^{t, x; u^{y_l}, v^{y_l}}) - \varepsilon] 1_{O_l}(X_{t_i}^{t, x; u^\varepsilon, v^\varepsilon}) \\ &= \sum_{l \geq 1} W_j(t_i, X_{t_i}^{t, x; u^{y_l}, v^{y_l}}) 1_{O_l}(X_{t_i}^{t, x; u^\varepsilon, v^\varepsilon}) - \varepsilon \\ &= W_j(t_i, X_{t_i}^{t, x; u^\varepsilon, v^\varepsilon}) - \varepsilon, \end{aligned}$$

from which we get the desired result. $\square$

Let us come to the proof of Proposition 5.6.

Proof. Let $t = t_0 < t_1 < \dots < t_n = T$ be a partition of $[t, T]$, and $\tau = \sup_i (t_{i+1} - t_i)$. By Proposition 3.9 and Lemma 5.4 we see that, for all $j = 1, 2$, $0 \leq k \leq n$, $s \in [t_k, t_{k+1})$ and $(u, v) \in \mathcal{U}_{t, T} \times \mathcal{V}_{t, T}$,

$$\begin{aligned} & \mathbb{E}[|W_j(t_k, X_{t_k}^{t, x; u, v}) - W_j(s, X_s^{t, x; u, v})|^2] \\ &\leq 2\mathbb{E}[|W_j(t_k, X_{t_k}^{t, x; u, v}) - W_j(s, X_{t_k}^{t, x; u, v})|^2] \\ &\quad + 2\mathbb{E}[|W_j(s, X_{t_k}^{t, x; u, v}) - W_j(s, X_s^{t, x; u, v})|^2] \\ &\leq C|s - t_k|(1 + \mathbb{E}[|X_{t_k}^{t, x; u, v}|^2]) + C\mathbb{E}[|X_{t_k}^{t, x; u, v} - X_s^{t, x; u, v}|^2] \\ &\leq C\tau. \end{aligned} \quad (5.26)$$

Here and after C represents a generic constant which may be different at different places.

We let Lemma 5.8 we let $(u^\varepsilon, v^\varepsilon) \in \mathcal{U}_{t, T} \times \mathcal{V}_{t, T}$ be defined as in Lemma 5.8 for $\varepsilon = \varepsilon_0$, where $\varepsilon_0 > 0$ will be specified later. Then, for all $i, 0 \leq i \leq n$,

$$W_j(t_i, X_{t_i}^{t, x; u^\varepsilon, v^\varepsilon}) - \varepsilon_0 \leq {}^j G_{t_i, t_{i+1}}^{t, x; u^\varepsilon, v^\varepsilon} [W_j(t_{i+1}, X_{t_{i+1}}^{t, x; u^\varepsilon, v^\varepsilon})], \mathbb{P} - a.s.$$

For $t \leq s_1 \leq s_2 \leq T$, let us suppose, without loss of generality, that $t_{i-1} \leq s_1 < t_i$ and $t_k < s_2 \leq t_{k+1}$, for some $1 \leq i < k \leq n-1$. Therefore, applying the Lemmas 2.3 and 2.4 we deduce that

$$\begin{aligned}
& {}^j G_{t_i, t_{k+1}}^{t, x; u^\varepsilon, v^\varepsilon} [W_j(t_{k+1}, X_{t_{k+1}}^{t, x; u^\varepsilon, v^\varepsilon})] \\
&= {}^j G_{t_i, t_k}^{t, x; u^\varepsilon, v^\varepsilon} [{}^j G_{t_k, t_{k+1}}^{t, x; u^\varepsilon, v^\varepsilon} [W_j(t_{k+1}, X_{t_{k+1}}^{t, x; u^\varepsilon, v^\varepsilon})]] \\
&\geq {}^j G_{t_i, t_k}^{t, x; u^\varepsilon, v^\varepsilon} [W_j(t_k, X_{t_k}^{t, x; u^\varepsilon, v^\varepsilon}) - \varepsilon_0] \\
&\geq {}^j G_{t_i, t_k}^{t, x; u^\varepsilon, v^\varepsilon} [W_j(t_k, X_{t_k}^{t, x; u^\varepsilon, v^\varepsilon})] - C\varepsilon_0 \\
&\geq \dots \geq {}^j G_{t_i, t_{i+1}}^{t, x; u^\varepsilon, v^\varepsilon} [W_j(t_{i+1}, X_{t_{i+1}}^{t, x; u^\varepsilon, v^\varepsilon})] - C(k-i)\varepsilon_0 \\
&\geq W_j(t_i, X_{t_i}^{t, x; u^\varepsilon, v^\varepsilon}) - C(k-i+1)\varepsilon_0,
\end{aligned}$$

from which we see that

$$\begin{aligned}
& {}^j G_{s_1, t_{k+1}}^{t, x; u^\varepsilon, v^\varepsilon} [W_j(t_{k+1}, X_{t_{k+1}}^{t, x; u^\varepsilon, v^\varepsilon})] \\
&= {}^j G_{s_1, t_i}^{t, x; u^\varepsilon, v^\varepsilon} [{}^j G_{t_i, t_{k+1}}^{t, x; u^\varepsilon, v^\varepsilon} [W_j(t_{k+1}, X_{t_{k+1}}^{t, x; u^\varepsilon, v^\varepsilon})]] \\
&\geq {}^j G_{s_1, t_i}^{t, x; u^\varepsilon, v^\varepsilon} [W_j(t_i, X_{t_i}^{t, x; u^\varepsilon, v^\varepsilon}) - C(k-i+1)\varepsilon_0] \\
&\geq {}^j G_{s_1, t_i}^{t, x; u^\varepsilon, v^\varepsilon} [W_j(t_i, X_{t_i}^{t, x; u^\varepsilon, v^\varepsilon})] - C(k-i+2)\varepsilon_0 \\
&\geq {}^j G_{s_1, t_i}^{t, x; u^\varepsilon, v^\varepsilon} [W_j(t_i, X_{t_i}^{t, x; u^\varepsilon, v^\varepsilon})] - \frac{\varepsilon}{2},
\end{aligned}$$

where we put $\varepsilon_0 = \frac{\varepsilon}{2Cn}$. We set

$$\begin{aligned}
I_1 &= {}^j G_{s_1, t_{k+1}}^{t, x; u^\varepsilon, v^\varepsilon} [W_j(t_{k+1}, X_{t_{k+1}}^{t, x; u^\varepsilon, v^\varepsilon})] - {}^j G_{s_1, t_i}^{t, x; u^\varepsilon, v^\varepsilon} [W_j(t_i, X_{t_i}^{t, x; u^\varepsilon, v^\varepsilon})] + \frac{\varepsilon}{2} \geq 0, \\
I_2 &= {}^j G_{s_1, s_2}^{t, x; u^\varepsilon, v^\varepsilon} [W_j(s_2, X_{s_2}^{t, x; u^\varepsilon, v^\varepsilon})] - W_j(s_1, X_{s_1}^{t, x; u^\varepsilon, v^\varepsilon}) + \frac{\varepsilon}{2}.
\end{aligned} \tag{5.27}$$

In what follows we shall prove the following:

$$\mathbb{E}[|I_1 - I_2|^2] \leq C\tau.$$

Let us put

$$y_s = {}^j G_{s, t_i}^{t, x; u^\varepsilon, v^\varepsilon} [W_j(t_i, X_{t_i}^{t, x; u^\varepsilon, v^\varepsilon})], s \in [s_1, t_i].$$

Then we consider the associated BSDEs:

$$\left\{ \begin{array}{l} y_s = W_j(t_i, X_{t_i}^{t, x; u^\varepsilon, v^\varepsilon}) + \int_s^{t_i} f_j(r, X_r^{t, x; u^\varepsilon, v^\varepsilon}, y_r, z_r, u_r^\varepsilon, v_r^\varepsilon) dr + k_{t_i} - k_s - \int_s^{t_i} z_r dB_r, \\ y_s \geq h_j(s, X_s^{t, x; u^\varepsilon, v^\varepsilon}), \quad s \in [s_1, t_i], \\ {}^2K_{t_i} = 0, \quad \int_{t_i}^{t_{i+1}} (y_r - h_j(r, X_r^{t, x; u^\varepsilon, v^\varepsilon})) dk_r = 0, \end{array} \right.$$

and

$$y'_s = W_j(s_1, X_{s_1}^{t, x; u^\varepsilon, v^\varepsilon}), \quad s \in [s_1, t_i].$$

Then applying Lemma 2.3 we conclude

$$\begin{aligned}
& |{}^j G_{s_1, t_i}^{t, x; u^\varepsilon, v^\varepsilon} [W_j(t_i, X_{t_i}^{t, x; u^\varepsilon, v^\varepsilon})] - W_j(s_1, X_{s_1}^{t, x; u^\varepsilon, v^\varepsilon})|^2 \\
&\leq C\mathbb{E}[|W_j(t_i, X_{t_i}^{t, x; u^\varepsilon, v^\varepsilon}) - W_j(s_1, X_{s_1}^{t, x; u^\varepsilon, v^\varepsilon})|^2 | \mathcal{F}_{s_1}] \\
&\quad + C\mathbb{E}[\int_{s_1}^{t_i} |f_j(r, X_r^{t, x; u^\varepsilon, v^\varepsilon}, y_r, z_r, u_r^\varepsilon, v_r^\varepsilon)|^2 | \mathcal{F}_{s_1}] \\
&\quad + C\mathbb{E}[\sup_{s_1 \leq s \leq t_i} |h_j(s, X_s^{t, x; u^\varepsilon, v^\varepsilon}) - h_j(s_1, X_{s_1}^{t, x; u^\varepsilon, v^\varepsilon})|^2 dr | \mathcal{F}_{s_1}]^{\frac{1}{2}}
\end{aligned}$$

$$\begin{aligned}
&\leq C\mathbb{E}[|W_j(t_i, X_{t_i}^{t,x;u^\varepsilon,v^\varepsilon}) - W_j(s_1, X_{s_1}^{t,x;u^\varepsilon,v^\varepsilon})|^2 | \mathcal{F}_{s_1}] \\
&\quad + C(t_i - s_1) + C\mathbb{E}[\sup_{s_1 \leq s \leq t_i} |X_s^{t,x;u^\varepsilon,v^\varepsilon} - X_{s_1}^{t,x;u^\varepsilon,v^\varepsilon}|^2 | \mathcal{F}_{s_1}]^{\frac{1}{2}},
\end{aligned}$$

where we have used assumptions (H3.3) and (H3.4) and the boundedness of f_j . Since $(u^\varepsilon, v^\varepsilon) \in \mathcal{U}_{t,T} \times \mathcal{V}_{t,T}$ is independent of $\mathcal{F}_t$ we have

$$\begin{aligned}
&\mathbb{E}[|{}^jG_{s_1,t_i}^{t,x;u^\varepsilon,v^\varepsilon}[W_j(t_i, X_{t_i}^{t,x;u^\varepsilon,v^\varepsilon})] - W_j(s_1, X_{s_1}^{t,x;u^\varepsilon,v^\varepsilon})|^2 | \mathcal{F}_t] \\
&\leq C\mathbb{E}[|W_j(t_i, X_{t_i}^{t,x;u^\varepsilon,v^\varepsilon}) - W_j(s_1, X_{s_1}^{t,x;u^\varepsilon,v^\varepsilon})|^2] + C(t_i - s_1) \\
&\quad + C\mathbb{E}[\sup_{s_1 \leq s \leq t_i} |X_s^{t,x;u^\varepsilon,v^\varepsilon} - X_{s_1}^{t,x;u^\varepsilon,v^\varepsilon}|^2 | \mathcal{F}_t]^{\frac{1}{2}}.
\end{aligned}$$

By virtue of (5.26) we have

$$\mathbb{E}[|{}^jG_{s_1,t_i}^{t,x;u^\varepsilon,v^\varepsilon}[W_j(t_i, X_{t_i}^{t,x;u^\varepsilon,v^\varepsilon})] - W_j(s_1, X_{s_1}^{t,x;u^\varepsilon,v^\varepsilon})|^2] \leq C\tau^{\frac{1}{2}}. \quad (5.28)$$

By a similar argument

$$\mathbb{E}[|{}^jG_{s_2,t_{k+1}}^{t,x;u^\varepsilon,v^\varepsilon}[W_j(t_{k+1}, X_{t_{k+1}}^{t,x;u^\varepsilon,v^\varepsilon})] - W_j(s_2, X_{s_2}^{t,x;u^\varepsilon,v^\varepsilon})|^2] \leq C\tau^{\frac{1}{2}}. \quad (5.29)$$

For $s \in [s_1, s_2]$ we put

$$y_s^1 = {}^jG_{s,t_{k+1}}^{t,x;u^\varepsilon,v^\varepsilon}[W_j(t_{k+1}, X_{t_{k+1}}^{t,x;u^\varepsilon,v^\varepsilon})] = {}^jG_{s,s_2}^{t,x;u^\varepsilon,v^\varepsilon}[{}^jG_{s_2,t_{k+1}}^{t,x;u^\varepsilon,v^\varepsilon}[W_j(t_{k+1}, X_{t_{k+1}}^{t,x;u^\varepsilon,v^\varepsilon})]],$$

and

$$y_s^2 = {}^jG_{s,s_2}^{t,x;u^\varepsilon,v^\varepsilon}[W_j(s_2, X_{s_2}^{t,x;u^\varepsilon,v^\varepsilon})].$$

Let us consider the associated BSDEs:

$$\left\{ \begin{array}{l} y_s^1 = {}^jG_{s_2,t_{k+1}}^{t,x;u^\varepsilon,v^\varepsilon}[W_j(t_{k+1}, X_{t_{k+1}}^{t,x;u^\varepsilon,v^\varepsilon})] + \int_s^{s_2} f_j(r, X_r^{t,x;u^\varepsilon,v^\varepsilon}, y_r^1, z_r^1, u_r^\varepsilon, v_r^\varepsilon) dr \\ \quad + k_{s_2}^1 - k_s^1 - \int_s^{s_2} z_r^1 dB_r, \\ y_s^1 \geq h_j(s, X_s^{t,x;u^\varepsilon,v^\varepsilon}), \quad s \in [s_1, s_2], \\ k_{s_2}^1 = 0, \quad \int_{s_1}^{s_2} (y_r - h_j(r, X_r^{t,x;u^\varepsilon,v^\varepsilon})) dk_r^1 = 0, \end{array} \right.$$

and

$$\left\{ \begin{array}{l} y_s^2 = W_j(s_2, X_{s_2}^{t,x;u^\varepsilon,v^\varepsilon}) + \int_s^{s_2} f_j(r, X_r^{t,x;u^\varepsilon,v^\varepsilon}, y_r^2, z_r^2, u_r^\varepsilon, v_r^\varepsilon) dr \\ \quad + k_{s_2}^1 - k_s^1 - \int_s^{s_2} z_r^2 dB_r, \\ y_s^2 \geq h_j(s, X_s^{t,x;u^\varepsilon,v^\varepsilon}), \quad s \in [s_1, s_2], \\ k_{s_2}^2 = 0, \quad \int_{s_1}^{s_2} (y_r - h_j(r, X_r^{t,x;u^\varepsilon,v^\varepsilon})) dk_r^2 = 0, \end{array} \right.$$

From the Lemmas 2.3 and 3.1 it follows that

$$\begin{aligned}
&|{}^jG_{s_1,t_{k+1}}^{t,x;u^\varepsilon,v^\varepsilon}[W_j(t_{k+1}, X_{t_{k+1}}^{t,x;u^\varepsilon,v^\varepsilon})] - {}^jG_{s_1,s_2}^{t,x;u^\varepsilon,v^\varepsilon}[W_j(s_2, X_{s_2}^{t,x;u^\varepsilon,v^\varepsilon})]|^2 \\
&\leq C\mathbb{E}[|{}^jG_{s_2,t_{k+1}}^{t,x;u^\varepsilon,v^\varepsilon}[W_j(t_{k+1}, X_{t_{k+1}}^{t,x;u^\varepsilon,v^\varepsilon})] - W_j(s_2, X_{s_2}^{t,x;u^\varepsilon,v^\varepsilon})|^2 | \mathcal{F}_{s_1}].
\end{aligned}$$

Hence, by (5.29) we see that

$$\mathbb{E}[|{}^jG_{s_1,t_{k+1}}^{t,x;u^\varepsilon,v^\varepsilon}[W_j(t_{k+1}, X_{t_{k+1}}^{t,x;u^\varepsilon,v^\varepsilon})] - {}^jG_{s_1,s_2}^{t,x;u^\varepsilon,v^\varepsilon}[W_j(s_2, X_{s_2}^{t,x;u^\varepsilon,v^\varepsilon})]|^2] \leq C\tau^{\frac{1}{2}}.$$

The above inequality and (5.28) yield

$$\mathbb{E}[|I_1 - I_2|^2] \leq C\tau^{\frac{1}{2}}.$$

Therefore,

$$\mathbb{P}(I_2 \leq -\frac{\varepsilon}{2}) \leq \mathbb{P}(|I_1 - I_2| \geq \frac{\varepsilon}{2}) \leq \frac{4\mathbb{E}[|I_1 - I_2|^2]}{\varepsilon^2} \leq \frac{4C\tau^{\frac{1}{2}}}{\varepsilon^2} \leq \varepsilon,$$

where we choose $\tau \leq \left(\frac{\varepsilon^3}{4C}\right)^2$, and by (5.27) we have

$$\mathbb{P}\left(W_j(s_1, X_{s_1}^{t,x;u^\varepsilon,v^\varepsilon}) - \varepsilon \leq {}^jG_{s_1,s_2}^{t,x;u^\varepsilon,v^\varepsilon}[W_j(s_2, X_{s_2}^{t,x;u^\varepsilon,v^\varepsilon})]\right) \geq 1 - \varepsilon.$$

We also refer to the fact that since $(u^\varepsilon, v^\varepsilon)$ is independent of $\mathcal{F}_t$, the conditional probability $\mathbb{P}(\cdot|\mathcal{F}_t)$ of the event $\left\{W_j(s_1, X_{s_1}^{t,x;u^\varepsilon,v^\varepsilon}) - \varepsilon \leq {}^jG_{s_1,s_2}^{t,x;u^\varepsilon,v^\varepsilon}[W_j(s_2, X_{s_2}^{t,x;u^\varepsilon,v^\varepsilon})]\right\}$ coincides with its probability. Indeed, also $\left\{W_j(s_1, X_{s_1}^{t,x;u^\varepsilon,v^\varepsilon}) - \varepsilon \leq {}^jG_{s_1,s_2}^{t,x;u^\varepsilon,v^\varepsilon}[W_j(s_2, X_{s_2}^{t,x;u^\varepsilon,v^\varepsilon})]\right\}$ is independent of $\mathcal{F}_t$. The proof is complete. $\square$

Finally, we give another main result: the existence theorem of a Nash equilibrium payoff.

Theorem 5.9. *Under the Isaacs condition (4.10), for all $(t, x) \in [0, T] \times \mathbb{R}^n$, there exists a Nash equilibrium payoff at (t, x) .*

Proof. By Theorem 5.5 we only have to prove that, for all $\varepsilon > 0$, there exists $(u^\varepsilon, v^\varepsilon) \in \mathcal{U}_{t,T} \times \mathcal{V}_{t,T}$ which satisfies (5.4) and (5.5) for $s \in [t, T], j = 1, 2$. For $\varepsilon > 0$, let us consider $(u^\varepsilon, v^\varepsilon) \in \mathcal{U}_{t,T} \times \mathcal{V}_{t,T}$ given by Proposition 5.6, i.e., in particular, $(u^\varepsilon, v^\varepsilon)$ is independent of $\mathcal{F}_t$. Setting $s_1 = t$ and $s_2 = T$ in Proposition 5.6, we get (5.4). Since $(u^\varepsilon, v^\varepsilon)$ is independent of $\mathcal{F}_t$, $J_j(t, x; u^\varepsilon, v^\varepsilon), j = 1, 2$, are deterministic and $\left\{(J_1(t, x; u^\varepsilon, v^\varepsilon), J_2(t, x; u^\varepsilon, v^\varepsilon)), \varepsilon > 0\right\}$ is a bounded sequence. Therefore, we can choose an accumulation point of this sequence, as $\varepsilon \rightarrow 0$. We denote this point by (e_1, e_2) . From Theorem 5.5 we see that (e_1, e_2) is a Nash equilibrium payoff at (t, x) . The proof is complete. $\square$

Acknowledgements.

The author thanks Prof. Rainer Buckdahn for his suggestions and comments.

References

- [1] Fleming, W.H., Souganidis, P.E.: On the existence of value functions of twoplayer, zero-sum stochastic differential games. Indiana Univ. Math. J. 38, 293–314 (1989)
- [2] Buckdahn, R., Li, J.: Stochastic differential games and viscosity solutions of Hamilton-Jacobi-Bellman-Isaacs equations. SIAM J. Control Optim. 47 444–475 (2008)
- [3] Buckdahn, R., Cardaliaguet, P., Quincampoix, M.: Some recent aspects of differential game theory. Dynamic Games and Applications 1, 74–114 (2011)
- [4] El Karoui, N., Kapoudjian, C., Pardoux, E., Peng, S., Quenez, M.C.: Reflected solutions of backward SDE's, and related obstacle problems for PDE's. Annals of Probability 25, 702–737 (1997)
- [5] El Karoui, N., Peng, S., Quenez, M.C.: Backward stochastic differential equation in finance. Math. Finance 7 1–71 (1997)
- [6] Wu, Z. Yu, Z.: Dynamic programming principle for one kind of stochastic recursive optimal control problem and Hamilton-Jacobi-Bellman equation. SIAM J. Control Optim. 47, 2616–2641 (2008)

- [7] Buckdahn, R., Cardaliaguet, P., Rainer, C.: Nash equilibrium payoffs for nonzero-sum Stochastic differential games. *SIAM J. Control Optim.* 43, 624–642 (2004)
- [8] Lin, Q.: A BSDE approach to Nash equilibrium payoffs for stochastic differential games with non-linear cost functionals. *Stochastic Processes and their Applications*, doi :10.1016/j.spa.2011.08.011.
- [9] Lin, Q.: Nash equilibrium payoffs for stochastic differential games with jumps and coupled nonlinear cost functionals, arXiv:1108.3695v1.
- [10] Lin, Q.: Nash equilibrium payoffs for N players stochastic differential games, preprint.
- [11] Lin, Q.: Nash equilibrium payoffs for two-player nonzero-sum stochastic differential games with two reflecting obstacles, preprint.
- [12] Lin, Q.: Stochastic optimal control problems of fully coupled forward backward stochastic differential equations with jumps, preprint.
- [13] Buckdahn, R., Li, J.: Stochastic differential games with reflection and related obstacle problems for Isaacs equations, arxiv0707.1133.
- [14] Hamadène, S., Lepeltier, J., Matoussi, A.: Double barrier backward SDEs with continuous coefficient. In: El Karoui Mazliak (Ed.): *Backward Stochastic Differential Equations*, Pitman Research Notes in Math. Series, vol. 364, 1997, pp. 161–175.
- [15] Peng, S.: Backward stochastic differential equations—stochastic optimization theory and viscosity solutions of HJB equations. In: Yan, J., Peng, S., Fang, S., Wu, L.M.: *Topics in Stochastic Analysis*, Ch.2 (Chinese vers.), (1997).
- [16] Crandall, M.G., Ishii, H., Lions, P.L.: User’s guide to viscosity solutions of second order partial differential equations. *Bull. Amer. Soc.* 27 1–67 (1992)
- [17] Barles, G., Buckdahn R., Pardoux, E.: Backward stochastic differential equations and integral-partial differential equations. *Stochastics* 60, 57–83 (1997)
- [18] Hamadène, S.: Backward-forward SDEs and stochastic differential games. *Stochastic Process. Appl.* 77, 1–15 (1998)